

similans Becker, 1924, Ent. Mitt. 13: 15 (δ ♀; *Pipunculus*). Type-loc: Taihoku, Formosa. For complete synonymy refer to Hardy (1956: 4).

sylvatica (Meigen). — Widespread Oriental; Palaearctic, Nearctic, Ethiopian regions.

1824, Syst. Beschr. europ. zweifl. Insekten. 4: 20 (sex?). Type-loc: Europe.

synadelpha Perkins. — Philippines [Luzon]; Australia, Bismarck Islands.

1905, Bull. Hawaiian Sug. Plrs' Ass. Exp. Sta 1(4): 150 (δ ; *Pipunculus*). Type-loc: Bundaberg, Queensland, Australia.

synadelphoides (de Meijere). — Java, ?Philippines [Luzon].

1914, Tijdschr. Ent. 57: 172 (δ ♀; *Pipunculus*). Type-loc: Semarang, Java.

unnamed sp. nr. *nyctias*. — Philippines [Luzon, Busuang].

Hardy, 1968, Ent. Meddr 36: 490 (δ).

unnamed sp. nr *subvirescens*. — Philippines [Tawi Tawi].

1968, Ent. Meddr 36: 489 (♀).

Unplaced species of Pipunculidae

gracilis Kertész. — Formosa.

1912, Annls hist.-nat. Mus. natn. hung. 10: 287 (δ ; *Dorylas*). Type-loc: Takao, Formosa. Probably *Dorylomorpha*.

mirabilis Brunetti. — India [Assam].

1912, Rec. Indian Mus. 7: 492 (♀; *Pipunculus*). Type-loc: Margherita, Assam, India. Probably *Tomosvaryella*.

Family SYRPHIDAE

By Lloyd V. Knutson

Systematic Entomology Laboratory, Entomology Research Division, ARS,
 USDA, c/o U.S. National Museum, Washington, D. C. 20560

F. Christian Thompson

10 Edmunds Road, Wellesley Hills, Massachusetts 02181

J. Richard Vockeroth

Biosystematics Research Institute, Agriculture Canada,
 Ottawa, Canada

The Syrphidae, "hover flies" or "flower flies," are one of the major groups of Diptera, with about 5,000 described species. The adults of many species are mimics of bees or wasps, are strikingly colored, or of unusual structure, and they are frequently collected. Many adults visit flowers and they are of great value in pollination of economically important plants. Adult Syrphidae are easily recognized by the special wing venation: spurious longitudinal vein present (except *Graptomyza*), basal cells long, and apical cells closed. Larvae and puparia can be distinguished by the contiguous posterior spiracular tubes. The larvae of many Syrphinae are important natural enemies of aphids, scale insects, and other small, soft-bodied insects. Larvae of Milesiinae are generally saprophagous in many kinds of aquatic and semi-aquatic situations. Larvae of Microdontinae are associated with ant nests. None of the Syrphidae are pests except for a few species of *Eristalis* that occasionally cause human myiasis and a few species that are important saprophages in bulbs of cultivated plants.

A total of 84 genera and 771 species are known from the Oriental region. All of the original literature was examined during preparation of this catalog. Included are 80 new combinations, 5 new generic, and 15 new specific synonyms; most of these are based on recent type examinations by J. R. Vockeroth. Five species are given new names.

We have followed the established usage of the generic names *Syrphus* Fabricius, *Cheilosia* sensu Meigen, *Eristalis* Latreille, *Xylota* Meigen, and *Milesia* Latreille. See Stone, et al. (1965) for a more complete synonymy and discussion of the nomenclature problems involved with these genera (except *Milesia*, see below).

We thank George C. Steyskal for checking the nomenclature and offering many improvements in grammatical treatment.

References: Bhatia 1931, 1939; Bhatia and Shaffi 1933 (biol., imm. stages Indian spp.); Boyes, et al. 1964-1971 (cytology); Brunetti 1923b (India); Chandler 1968 (eggs); Coe 1964 (Nepal); Curran 1928, 1931a, b (Malaya, Borneo); Dixon 1960 (imm. stages British spp.); Hartley 1963 (larval morph.); Heiss 1938 (class. imm. stages III. spp.); Hennig 1952 (world rev., imm. stages); Hull 1936 (checklist Indo-Austr.), 1949 (generic rev. for world); Keiser 1958 (Ceylon); Kertész 1910 (world cat.); Metcalf 1921 (δ genitalia); Roberts 1970 (larval morph.); Schneider 1969 (bionomics, physiol.); Shiraki 1930 (Formosa).

Subfamily SYRPHINAE

References: Schneider 1969 (biol. aphidophagous spp.); Vockeroth 1969 (tribal classif.).

Tribe SYRPHINI

Reference: Vockeroth 1969 (rev. world genera).

Genus ALLOGRAPTA Osten Sacken

Allograpta Osten Sacken, 1875, Bull. Buffalo Soc. nat. Sci. 3: 49. Type-species, *Scaeva obliqua* Say, by monotypy.

Miogramma Frey, 1946, Notul. ent. 25: 165. Type-species, *Syrphus javanus* Wiedemann, by original designation.

Helenomyia Bankowska, 1962, Bull. Acad. pol. Sci. Cl. II. Ser. Sci. Biol. 10: 311. Type-species, *Syrphus javanus* Wiedemann, by original designation.

References: Frey 1946 (key); Bhatia and Shaffi 1933 (biol., imm. stages).

javana (Wiedemann). — Borneo, Ceylon, China, Formosa, India [Madras, Mysore], Java, Malaya, Philippines, Sumatra, Thailand; Australia, Fiji, Guadalcanal, Hawaii, Japan, Korea, New Guinea.

1824, Analecta Ent.: 34 (sex?; *Syrphus*). Type-loc: Java.

distinctus Kertész, 1899, Termesztr. Füz. 22: 177 (♀; *Melithreptus*). Type-loc: Erima, Astrolabe Bay, New Guinea.

kinabaluensis (Curran). — Borneo.

1931, J. fed. Malay St. Mus. 16: 350 (♂ ♀; *Syrphus*). Type-loc: Mt Kinabalu, Marei Parei (5000 ft), N Borneo.

maculipleura (Brunetti). — India [Assam, Madras], Laos.

1913, Rec. Indian Mus. 8: 162 (♀; *Syrphus*). Type-loc: Rotung, NE Frontier, Assam, India.

bouvieri Hervé-Bazin, 1923, Bull. Soc. ent. Fr. 1923: 26 (♂ ♀; *Xanthogramma*). Type-locs: Laos; Trichinopoly and Coonoor, Nilghiri, India.

medanensis (de Meijere). — Sumatra.

1914, Tijdschr. Ent. 57: 165 (♂ ♀; as *Sphaerophoria javana* var.). Type-loc: Medan, Sumatra.

nasutus (Macquart). — Not Oriental; Ethiopian.

obscuricornis (de Meijere). — Java.

1914, Tijdschr. Ent. 57: 165 (♂; *Sphaerophoria*). Type-loc: Gunung Gede, Java.

philippina (Frey). — Philippines [Luzon].

1946, Notul. ent. 25: 167 (♀; *Miogramma*). Type-loc: Balbalasang, Luzon, Philippines.

purpureicollis (Frey). — Sumatra.

1946, Notul. ent. 25: 167 (♂; *Miogramma*). Type-loc: Ost-Sumatra.

Syrphidae

robinsoni (Curran). — Malaya.

1928, J. fed. Malay St. Mus. 14: 208 (♂ ♀; *Syrphus*). Type-loc: Rhododendron Hill, Cameron Highlands (5200 ft), Pahang, Malaya.

nigrotibialis Curran, 1928, J. fed. Malay St. Mus. 14: 241 (♀; *Sphaerophoria*). Type-loc: Kuala Tahan (300 ft), Pahang, Malaya. N. synonymy..

Genus ASARKINA Macquart

Asarkina Macquart, 1842, Dipt. exot. 2(2): 77 (137). Type-species, *Scaeva rostrata* Wiedemann, by monotypy.

Ancylosyrphus Bigot, 1882, Annls Soc. ent. Fr. (6) 2 (Bull): lxviii. Type-species, *Syrphus salviae* Fabricius, by original designation.

Reference: Curran 1931 (key).

Subgenus ASARKINA Macquart

assimilis (Macquart). — India.

1846, Dipt. exot. Suppl. 1: 135 (263) (♂; *Syrphus*). Type-loc: India. N. comb.

biroi Bezz. — Borneo, Java, Malaya, Sumatra, Sumbawa; New Guinea.

1908, Annls hist.-nat. Mus. natn. hung. 6: 502 (♀). Type-loc: Erima, Astrolabe Bay, New Guinea.

cingulata (Bigot). — Sula; Moluccas (Mysol).

1884, Annls Soc. ent. Fr. (6) 4: 85 (♀; *Melanostoma*). Type-loc: Mysol, Moluccas. N. comb.

consequens (Walker). — Borneo, Celebes, Formosa, India, Java, Malaya, Philippines [Samar, Palawan, Mindanao], Sumatra, Sumbawa; New Guinea, Samoa.

1857, J. Linn. Soc. Lond. 1: 18 (♂ ♀; *Syrphus*). Type-locs: Singapore, and M: Ophir (Malaya).

striata van der Wulp, 1881, Midden-Sumatra Exped., Dipt. 4: 32 (♂ ♀; *Syrphus*). Type-locs: Rawas; Palembang, Sumatra.

kelantanensis Bezz. 1908, Annls hist.-nat. Mus. natn. hung. 6: 499 (♀; as *ericetorum* var.). Type-loc: Kelantan, Malacca, Malaya.

var. *exilis* Curran. — Malaya.

1928, J. fed. Malay St. Mus. 14: 235 (♂). Type-loc: Jor Camp (1800 ft). Batang Padang, Perala, Malaya.

ericetorum (Fabricius). — throughout Oriental region; Africa, Australia, Fiji, Moluccas (Amboina), New Guinea, Samoa.

1781, Spec. Insect. 2: 425 (sex?; *Syrphus*). Type-loc: "Africa aequinoctial."

incisuralis Macquart, 1855, Dipt. exot. Suppl. 5: 94 (114) (♀; *Syrphus*). Type-loc: "Inde."

diaphana Doleschall, 1857, Natuurk. Tijdschr. Ned.-Indië 14: 409 (sex?; *Didea*). Type-loc: Amboina, Moluccas.

macquarti Doleschall, 1857, Natuurk. Tijdschr. Ned.-Indië 14: 408 (sex?; *Didea*). Type-loc: Amboina, Moluccas.

ssp. *formosae* Bezz. — Ceylon, Formosa, Sumbawa.

1908, Annls hist.-nat. Mus. natn. hung. 6: 499 (♂ ♀). Type-loc: Takao, Formosa.

- eurytaeniata* Bezz. — Bali, Borneo, Flores, Formosa, Malaya, Philippines [Luzon, Mindanao], Simeuluë Island, Sumatra, Guadalcanal.
 1908, Annls hist.-nat. Mus. natn. hung. 6: 501 (♀). Type-loc: Kelantan, Malacca, Malaya.
- fumipennis* Sack. — Formosa.
 1913, Ent. Mitt. 2: 3 (♀). Type-locs: Yamo, and Chip-Chip, Formosa.
- incompleta* Curran. — Java, Malaya.
 1928, J. fed. Malay St. Mus. 14: 239 (♂ ♀). Type-locs: Gunung Terbakar (4500 ft), Lubok Tamang, Pahang, Malaya; and Papandajan (5500–7000 ft), Preanger Regency, Java.
- laticornis* Curran. — Malaya, Thailand.
 1928, J. fed. Malay St. Mus. 14: 235 (♂). Type-loc: Khao Luang (5800 ft), Nakhon Si Thammarat, Thailand.
- morokaensis* de Meijere. — Borneo, Java; New Guinea.
 1908, Tijdschr. Ent. 51: 308 (♂ ♀). Type-loc: Moroka, New Guinea.
- nigripes* Curran. — Malaya.
 1923, J. fed. Malay St. Mus. 14: 236 (♂). Type-loc: Tanah Rata (4800 ft), Cameron Highlands, Pahang, Malaya.
- orientalis* Bezz. — China, Malaya, Philippines [Luzon, Palawan].
 1908, Annls hist.-nat. Mus. natn. hung. 6: 498 (♂ ♀; as *ericorum* var.). Type-locs: China, and Malaya.
- porcina* porcina (Coquillet). — Ceylon, China [Fukien], Formosa, India [Assam, W Bengal]; Japan.
 1898, Proc. U.S. natn. Mus. 21: 322 (♂ ♀; *Syrphus*). Type-loc: Japan.
- ribbei* Bezz. — Sumbawa; New Britain, New Guinea.
 1908, Annls hist.-nat. Mus. natn. hung. 6: 501 (♂ ♀). Type-locs: Kinigung, New Pomerania, and Friedrich-Wilhelmshafen, Stephansort, New Guinea.
- rufofasciata* (Macquart). — Java.
 1850, Dipt. exot. Suppl. 4: 149 (453) (♀; *Syrphus*). Type-loc: Java. N. comb.
- salviae* (Fabricius). — Borneo, India, Malaya; Africa, Guadalcanal, Moluccas (Amboina).
 1794, Ent. Syst. 4: 306 (sex?; *Syrphus*). Type-loc: Sierra Leone, Africa.

Genus BETASYRPHUS Matsumura

- Betasyrphus* Matsumura, 1917, Ent. Mag. Kyoto 2: 143. Type-species, *Syrphus serarius* Wiedemann, by original designation.
References: Bhatia and Shafii 1933 (biol., imm. stages).
- albibasis* (Coe). — Nepal.
 1964, Bull. Br. Mus. nat. Hist. (Ent.) 15: 263 (♀; *Syrphus*). Type-loc: Arun Valley, Tumlingtar Plateau, E Nepal. N. comb.
- serarius* (Wiedemann). — throughout SE Asia; Australia, Japan, Korea, New Guinea.
 1830, Aussereurop. zweifl. Insekts. 2: 128 (♀; *Syrphus*). Type-loc: China.

Genus CITROGRAMMA Vockeroth

- Citrogramma* Vockeroth, 1969, Mem. ent. Soc. Can. 62: 92. Type-species, *Syrphus herbebazini* Curran, by original designation.
- circumdatum* (de Meijere). — Borneo; New Guinea.
 1908, Tijdschr. Ent. 51: 306 (♂ ♀; *Syrphus*). Type-loc: Moroka, New Guinea. N. comb.
- citrinum* (Brunetti). — Ceylon, India [Assam, Mysore], Malaya, Thailand.
 1923, Fauna Br. India, Dipt. 3: 95 (♂ ♀; *Xanthogramma*). Type-loc: Mangaldai, Assam, India.
- clarum* (Hervé-Bazin). — Borneo, Java, Laos, Philippines [Luzon].
 1923, Bull. Soc. ent. Fr. 1923: 25 (♂; *Oltiosyrphus*). Type-loc: Chapa (Tonkin), Laos.
- difficile* (Curran). — Malaya.
 1928, J. fed. Malay St. Mus. 14: 213 (♂ ♀; *Syrphus*). Type-locs: Rhododendron Hill (5200 ft), and Cameron Highlands No. 4 Camp (4800 ft), Pahang, Malaya.
- fascipleura* (Curran). — Borneo.
 1931, J. fed. Malay St. Mus. 16: 352 (♂; *Syrphus*). Type-loc: Mt Kinabalu, Kenokok (3300 ft), N Borneo. N. comb.
- gedehanum* (de Meijere). — Java.
 1914, Tijdschr. Ent. 57: 156 (♂; *Syrphus*). Type-loc: Gunung Gede, Java. N. comb.
- herbebazini* (Curran). — Malaya; Sarawak.
 1928, J. fed. Malay St. Mus. 14: 207 (♂; *Syrphus*). Type-loc: Rhododendron Hill, Cameron Highlands (5200 ft), Pahang, Malaya.
- pendleburyi* (Curran). — Malaya.
 1928, J. fed. Malay St. Mus. 14: 206 (♂; *Syrphus*). Type-loc: Seat Point, Gunong Tahan, Pahang (5460 ft), Malaya. N. comb.
- variscutatum* (Curran). — Malaya, Thailand.
 1928, J. fed. Malay St. Mus. 14: 210 (♂ ♀; *Syrphus*). Type-locs: Seat Point (5460 ft), Gunong Tahan, Pahang (5500 ft); Padang (6000–7100 ft); Gunong Kechau (6300 ft); Cameron Highlands, Gunong Berumban (6050 ft); Rhododendron Hill (5200 ft), Pahang, Malaya, and Nakhon Si Thammarat, Khao Luang (2000 ft), Thailand.

Genus DASYSYRPHUS Enderlein

- Dasydysyrphus* Enderlein, 1938, Sber. Ges. naturf. Freunde Erl. 1937: 208. Type-species, *Scava albostriata* Fallén, by original designation.
- albostriatus* (Fallén). — Not Oriental; Palaearctic.
- bilineatus* (Matsumura). — Formosa; Japan.
 1917, Ent. Mag. Kyoto 3: 38 (♂; *Syrphus*). Type-loc: Sapporo, Hokkaido, Japan.
- brunettii* (Hervé-Bazin). — India [Himachal Pradesh, W Bengal], Nepal.
 1924, Annls Soc. ent. Fr. 92: 290 (*Syrphus*; n. name for *albostriatus* of Brunetti, not Fallén). N. comb.
- albostriatus* Brunetti, 1923, Fauna Br. India, Dipt. 3: 72 (♂ ♀; misidentification). Type-locs: Mundali, Dehra Dun District; Matiana, Simla District; Darjeeling and Kurseong, India.

- koningsbergeri* (de Meijere). — Borneo, Java.
 1914, Tijdschr. Ent. 57: 152 (♂; *Syrphus*). Type-loc: Pangerango, W Java.
- latistrigatus* (de Meijere). — Java.
 1914, Tijdschr. Ent. 57: 153 (♂; *Syrphus*). Type-loc: Pangerango, Java. N. comb.
- orsua* (Walker). — Ceylon, India, Sumatra.
 1852, Insecta Saundersiana 1: 231 (sex?; *Syrphus*). Type-loc: East Indies.
- serariooides* (de Meijere). — Borneo, Java.
 1924, Tijdschr. Ent. 67: 218 (♂ ♀; *Syrphus*). Type-loc: Pangerango, W Java.

Genus DIDEA Macquart

- Didea* Macquart, 1834, Hist. nat. Ins. Dipt. 1: 508. Type-species, *fasciata* Macquart, by monotypy.
- fasciata* Macquart. — Formosa, India [Assam]; Palaearctic.
 1834, Hist. nat. Ins. Dipt. 1: 508 (♂). Type-loc: Paris, France.

Genus DIDEOIDES Brunetti

- Dideoides* Brunetti, 1908, Rec. Indian Mus. 2: 54. Type-species, *ovatus* Brunetti, by monotypy.
- Dideodes* Matsumura, 1917, Ent. Mag. Kyoto 2: 140. Type-species, *Syrphus latus* Coquillett, by original designation.
- Malayomyia* Curran, 1928, J. fed. Malay St. Mus. 14: 225. Type-species, *pretiosus* Curran, by original designation.
- chrysotoxoides* (Curran). — Thailand.
 1928, J. fed. Malay St. Mus. 14: 201 (♀; *Syrphus*). Type-loc: Nakhon Si Thammarat, Khao Luang (5800 ft), Thailand. N. comb.
- consimilis* (Macquart). — Java.
 1849, Dipt. exot. Suppl. 4: 150 (454) (♂ ♀; *Syrphus*). Type-loc: Java. N. comb.
- coquilletti* (van der Goot). — China [Fukien], Formosa; Japan.
 1964, Beaufortia 10: 218 (*Syrphus*; n. name for *lautus* Coquillett).
- lautus* Coquillett, 1898, Proc. U.S. natn. Mus. 21: 322 (♀; *Syrphus*; preocc., Giglio-Tos 1892). Type-loc: Japan.
- depressus* (de Meijere). — Java, Sumatra.
 1914, Tijdschr. Ent. 57: 153 (♂; *Syrphus*). Type-loc: Gunung Gede, Java. N. comb.
- kempi* Brunetti. — India [Assam].
 1923, Fauna Br. India, Dipt. 3: 59 (♂ ♀). Type-loc: Tura, Garo Hills, Assam, India.
- latus* (Coquillett). — China [Fukien], Formosa; Japan.
 1898, Proc. U.S. natn. Mus. 21: 322 (♀; *Syrphus*). Type-loc: Japan.
- ssp. *formosanus* (Matsumura). — Formosa.
 1910, Keiseisha, Tokyo 46 (♂; *Syrphus*; as sp.). Type-loc: Taihoku, Tainan.
- ovatus* Brunetti. — India [Assam, Sikkim, W Bengal], Laos.
 1908, Rec. Indian Mus. 2: 54 (♂ ♀). Type-loc: Sikkim, India.

- pellucidipennis* (Coe). — Nepal.
 1964, Bull. Br. Mus. nat. Hist. (Ent.) 15: 259 (♀; *Syrphus*). Type-loc: Taplejung District, E Nepal. N. comb.
- pretiosus* (Curran). — Malaya.
 1928, J. fed. Malay St. Mus. 14: 226 (♂; *Malayomyia*). Type-loc: Rhododendron Hill (5200 ft), Cameron Highlands, Pahang, Malaya.
- tigerinus* (Bigot). — India [Assam].
 1885, Annls Soc. ent. Fr. (6) 5: 249 (♂; *Ischyrosyrphus*). Type-loc: India.
- trilineatus* Brunetti. — Burma.
 1923, Fauna Br. India, Dipt. 3: 62 (♂). Type-loc: Myitkyina District, Burma.

Genus DIDEOPSIS Matsumura

- Dideopsis* Matsumura, 1917, Ent. Ent. Mag. Kyoto 2: 142. Type-species, *Eristalis aegrotus* Fabricius, by original designation.
- Aegrotomyia* Frey, 1946, Notul. ent. 25: 158. Type-species, *Eristalis aegrotus* Fabricius, by original designation.
- aegrota* (Fabricius). — Nepal, throughout SE Asia; Australia, Moluccas (Amboina), New Guinea.
 1805, Syst. Antiat.: 243 (♀; *Eristalis*). Type-loc: China.
- scapipennis* Macquart, 1834, Hist. nat. Ins. Dipt. 1: 537 (♂; *Syrphus*). Type-loc: Java.
- ellenriederi* Doleschall, 1857, Natuurk. Tijdschr. Ned.-Indië 14: 407 (sex?; *Didea*). Type-loc: Amboina, Moluccas.
- infirmus* Rondani, 1875, Annali Mus. civ. Stor. nat. Giacomo Doria 7: 423 (♀; *Syrphus*). Type-loc: Sarawak, Borneo.
- pura* (Curran). — Indochina, Malaya.
 1928, J. fed. Malay St. Mus. 14: 230 (♀; *Asarkina*). Type-loc: Ampang (600 ft), Selangor, Malaya.

Genus EOSPHAEROPHORIA Frey

- Eosphaerophoria* Frey, 1946, Notul. ent. 25: 169. Type-species, *marginata* Frey, by original designation.
- Tambavanna* Keiser, 1958, Revue suisse Zool. 65: 202. Type-species, *dentiscutellata* Keiser, by original designation.
- dentiscutellata* (Keiser). — Ceylon, Indochina.
 1958, Revue suisse Zool. 65: 202 (♂; *Tambavanna*). Type-loc: Kandy, Deiyawella, Ceylon.
- marginata* Frey. — Philippines [Luzon].
 1946, Notul. ent. 25: 169 (♂ ♀). Type-loc: Mt Banahao, Luzon, Philippines.

Genus EPISTROPHE Walker

- Epistrophe* Walker, 1852, Insecta Saundersiana 3: 242. Type-species, *conjugens* Walker, by monotypy, = *grossulariae* (Meigen).

Subgenus EPISTROPHIE Walker

aequalis (Walker). — India [Punjab], ?Java.

1852, Insecta Saundersiana 1: 226 (♂; *Xylota*). Type-loc: East Indies. N. comb.

distinctus Brunetti, 1915, Rec. Indian Mus. 11: 211 (♂). Type-loc: Tolpan, Garhiwal District, W Himalaya, India.

exul (Curran). — Malaya.

1928, J. fed. Malay St. Mus. 14: 215 (♂ ♀; *Syrphus*). Type-locs: Rhododendron Hill (5200 ft), Cameron Highlands, and Seat Point (5460 ft), Gunong Tahan, Pahang, Malaya.

griseocincta (Brunetti). — India [Uttar Pradesh].

1923, Fauna Br. India, Dipt. 3: 77 (♂; *Syrphus*). Type-loc: Bogarkote, Kumaon, W Himalayas (8000 ft), India. N. comb.

hirsuteron (Curran). — Borneo.

1931, J. fed. Malay St. Mus. 16: 353 (♂; *Syrphus*). Type-loc: Mt Kinabalu, Pakka (10,200 ft), N Borneo.

Subgenus EPISTROPHELLA Dušek and Láska

Epistrophella Dušek and Láska, 1967, Přírodov. Pr. Česk. Akad. Věd. 1: 368. Type-species, *Syrphus euchromus* Kowarz, by original designation.

horishana (Matsumura). — Formosa, India [W Bengal], Laos, Philippines; Japan. 1917, Ent. Mag. Kyoto 3: 24 (♂; *Syrphus*). Type-loc: Philippines.

quinquevittatus Brunetti, 1923, Fauna Br. India, Dipt. 3: 81 (♂ ♀; *Syrphus*). Type-loc: Darjeeling District, Kalimpong, India.

Genus EPISYRPHUS Matsumura and Adachi

Episyphus Matsumura and Adachi, 1917, Ent. Mag. Kyoto 2: 134. Type-species, *Musca balteata* De Geer, by designation of Matsumura and Adachi (1917: 16). Reference: Bhatia and Shaffi 1933 (biol., imm. stages).

alternans (Macquart). — throughout Oriental region; Australia, Japan, Tibet. 1842, Dipt. exot. 2(2): 89 (149) (♂ ♀; *Syrphus*). Type-loc: Coromandel, India.

arcifer (Sack). — Formosa, Malaya, Nepal. 1927, Stettin. ent. Ztg 88: 306 (♂ ♀; *Syrphus*). Type-locs: Kankau, and Fuhosho, Formosa.

balteatus (De Geer). — entire Oriental region; Australia, Bonin Islands, Palaearctic.

1776, Mém. pour serv. Hist. Ins. 6: 116 (sex?; *Musca*). Type-loc: Europe. *nectareus* Fabricius, 1787, Mantissa Insect. 2: 341 (sex?; *Musca*). Type-loc: Denmark.

pleuralis Thonissen, 1869, in K. svenska fregatten Eugenies resa, Zool., Dipt.: 497 (♂; *Syrphus*). Type-loc: China.

triligatus Walker, 1857, J. Linn. Soc. Lond. 1: 19 (♂; *Syrphus*). Type-loc: Mt Ophir, Malaya.

contractus (Keiser). — Sumbawa. 1952, Verh. naturf. Ges. Basel 63: 159 (♀; *Epistrophe*). Type-loc: Sumbawa. N. comb.

Syrphidae

divertens (Walker). — Borneo, Formosa, Thailand.

1857, J. Linn. Soc. Lond. 1: 124 (♀; *Syrphus*). Type-loc: Sarawak, Borneo. *claviger* Sack, 1927, Stettin. ent. Ztg 88: 308 (♀; *Syrphus*). Type-loc: Fuhosho, Formosa.

graptus (Hull). — Formosa.

1944, Psyche, Camb. 51: 22 (♂; *Syrphus*). Type-loc: Sozan, Formosa.

heterogaster (Thomson). — China.

1869, in K. svenska fregatten Eugenies resa, Zool., Dipt.: 498 (♀; *Syrphus*). Type-loc: China. N. comb.

nectarinus (Wiedemann). — SE Asia to Japan; Australia, New Guinea.

1830, Aussereurop. zweifl. Insekt. 2: 128 (♂ ♀; *Syrphus*). Type-loc: China.

obligatus (Curran). — Malaya.

1931, J. fed. Malay St. Mus. 16: 316 (♂ ♀; *Syrphus*). Type-loc: Kedah Peak (3000–3950 ft), Malaya.

viridaureus (Wiedemann). — Java, Malaya; New Caledonia.

1824, Analecta Ent.: 35 (♂; *Syrphus*). Type-loc: Java.

Genus ERIOZONA Schiner

Eriozona Schiner, 1860, Wien. ent. Mschr. 4: 214. Type-species *Syrphus oestriformis* Meigen, by original designation, = *syrphoides* (Fallén).

analis Kertész. — India [Sikkim], Nepal.

1901, Természetr. Füz. 24: 414 (♂). Type-loc: Sikkim, India.

ruficauda Brunetti, 1907, Rec. Indian Mus. 1: pl. xi, fig. 2 (♀); 1908, Rec. Indian Mus. 2: 56. Type-loc: Sikkim, India.

nigroscutellata Shiraki. — Formosa.

1930, Mem. Fac. Agric. Taihoku imp. Univ. 1: 333 (♀). Type-locs: Royeichi, and Daisuikutsu, Formosa.

Genus ISCHIODON Sack

Ischiodon Sack, 1913, Ent. Mitt. 2: 5. Type-species, *trochanterica* Sack, by monotypy, = *scutellaris* (Fabricius).

References: Lal and Gupta 1953; Lal and Haque 1955 (biol., imm. stages).

scutellaris (Fabricius). — SE Asia to Japan; Australia, Hawaii, Micronesia. 1805, Syst. Antliat.: 252 (sex?; *Scaeva*). Type-loc: Tranquebar, India.

coromandensis Macquart, 1842, Dipt. exot. 2(2): 89 (149) (♂; *Syrphus*). Type-loc: côte de Coromandel (India).

splendens Doleschall, 1856, Natuurk. Tijdschr. Ned.-Indië 10: 410 (sex?; *Syrphus*). Type-loc: Java.

erythropygus Bigot, 1884, Annls Soc. ent. Fr. (6) 4: 87 (♂; *Syrphus*). Type-loc: "Indes."

novaeguineae Kertész, 1899, Természetr. Füz. 22: 178 (♂; *Melithreptus*). Type-locs: Friedrich-Wilhelmshafen and Erima, New Guinea.

trochanterica Sack, 1913, Ent. Mitt. 2: 6 (♂). Type-locs: Kanshizei, Polishe, Suihenkyaku, Tainah, and Takao, Formosa.

platychiroides Frey, 1946, Notul. ent. 25: 164 (♂). Type-loc: Atimonan, Quezon, Luzon, Philippines.

magnicornis Shiraki, 1963, Insects Micronesia 13: 141 (♀; *Epistrophe*). Type-loc:

Wena (Moen), Truk, Caroline Islands.

Genus ISCHYROSYRPHUS Bigot

Ischyrosyrphus Bigot, 1882, Annls Soc. ent. Fr. (6) 2 (Bull.): lxviii. Type-species, *sivae* Bigot, by designation of Kirby (1883: 244).
sivae Bigot. — India.
 1882, Annls Soc. ent. Fr. (6) 2 (Bull.): lxviii (♂). Type-loc: India.

Genus MELISCAEVA Frey

Epistrophe, subg. *Meliscaeva*, Frey, 1946, Notul. ent. 25: 164. Type-species, *Scaeva cinctella* Zetterstedt, by original designation.
 References: Scott 1939; Dixon 1960 (biol., imm. stages).
abdominalis (Sack). — Formosa.
 1927, Stettin. ent. Ztg 88: 310 (♂; *Baccha*). Type-loc: Toa Tsu Kutsu, Formosa. N. comb.
ceylonica (Keiser). — Ceylon.
 1958, Revue suisse Zool. 65: 198 (♂ ♀; *Metepistrophe*). Type-loc: Pidratalagala (2000–2400 m), Ceylon.
cinctella (Zetterstedt). — Ceylon, Formosa, India [Himachal Pradesh, W Bengal], Nepal; Palaearctic.
 1843, Dipt. Scandinaviae 2: 742 (♂ ♀; *Syrphus*). Type-loc: Europe.
var. formosana (Shiraki). — Formosa.
 1930, Mem. Fac. Agric. Taihoku imp. Univ. 1: 388 (♂ ♀; *Syrphus*). Type-locs: Shukoran, Royeichi, Daisukutsu, Arisan, Hassensan, Musha, Kanko, Yamanotaika, and Rantaisan, Formosa.
var. taiwana (Shiraki). — Formosa.
 1930, Mem. Fac. Agric. Taihoku imp. Univ. 1: 387 (♂ ♀; *Syrphus*). Type-locs: Musha and Arisan, Formosa.
deceptor (Curran). — Borneo, Malaya.
 1928, J. fed. Malay St. Mus. 14: 219 (♂; *Syrphus*). Type-locs: Gunong Kechau (6300 ft); Gunong Tahan, Padang Luas (4900 ft) and (6500 ft), Pahang, Malaya.
ichthyops (de Meijere). — Java.
 1914, Tijdschr. Ent. 57: 157 (♂ ♀; *Syrphus*). Type-loc: Pangerango, Gunung Gede, Java. N. comb.
malayensis (Curran). — Borneo, Malaya.
 1928, J. fed. Malay St. Mus. 14: 224 (♂; *Syrphus*). Type-locs: Rhododendron Hill (5200 ft) and Gunong Berumban (6050 ft), Cameron Highlands, Pahang, Malaya.
melanostomoides (Hull). — Java, Philippines [Mindanao].
 1941, Psyche, Camb. 48: 149 (♂ ♀; *Syrphus*). Type-loc: Tjibodas, Mt Gede (9000 ft), Java.
monticola (de Meijere). — Ceylon, Formosa, Java.
 1914, Tijdschr. Ent. 57: 159 (♀; *Syrphus*). Type-locs: Gunung Gede, and Tosari, Java. N. comb.
morna (Curran). — Borneo, Java, Malaya.
 1931, J. fed. Malay St. Mus. 16: 317 (♀; *Syrphus*). Type-loc: Fraser's Hill (4200 ft), Pahang, Malaya.

nigripes (de Meijere). — Java.

1914, Tijdschr. Ent. 57: 162 (♀; *Chamaesyrphus*). Type-loc: Gunung Gede, Java. N. comb.

peteus (Curran). — Borneo.

1931, J. fed. Malay St. Mus. 16: 349 (♀; *Syrphus*). Type-loc: Mt Kinabalu, Lumu Lumu (5500 ft), N Borneo. N. comb.

strigifrons (de Meijere). — Formosa, Jaya, Malaya.

1914, Tijdschr. Ent. 57: 158 (♂ ♀; as *Syrphus cinctellus* var.). Type-locs: Gede, and Nongkodjadjar, Java.

tenuiformis (Curran). — Borneo, Malaya.

1928, J. fed. Malay St. Mus. 14: 222 (♂; *Syrphus*). Type-loc: Gunong Tahan (6500–7100 ft), Pahang, Malaya.

Genus METASYRPHUS Matsumura

Metasyrphus Matsumura, 1917, Ent. Mag. Kyoto 2: 147. Type-species, *Scaeva corollae* Fabricius, by original designation.

References: Bhatia and Shaffi 1933; Bhatia 1939; Scott 1939 (biol., imm. stages).

Subgenus METASYRPHUS Matsumura

confrater (Wiedemann). — entire Oriental region; Japan, New Guinea.

1830, Aussereurop zweifl. Insek. 2: 120 (♀; *Syrphus*). Type-loc: China.
cranipes Walker, 1852, Insecta Saundersiana 1: 231 (♂ ♀; *Syrphus*). Type-loc: East Indies.

mundus Walker, 1852, Insecta Saundersiana 1: 230 (♂; *Syrphus*). Type-loc: East Indies.

trilimbatus Bigot, 1884, Annls Soc. ent. Fr. (6) 4: 86 (♂; *Syrphus*). Type-loc: "Indies."

corollae (Fabricius). — Asia including Formosa; N Africa, Palaearctic.
 1794, Ent. Syst. 4: 306 (sex?; *Scaeva*). Type-loc: "Killiae."

latifasciatus (Macquart). — India [Himachal Pradesh].

1829, Ins. Dipt. N. Fr. 4: 94 (242) (♂; *Syrphus*). Type-loc: Arras, France.

luniger (Meigen). — India [Punjab]; Canary Islands, Europe, Japan.

1822, Syst. Beschr. europ. zweifl. Insek. 3: 300 (♂; *Syrphus*). Type-loc: Europe.

macropterus (Thomson). — China.

1869, in K. svenska fregatten Eugenies resa, Zool., Dipt.: 498 (♀; *Syrphus*). Type-loc: China. N. comb.

torvooides (de Meijere). — Java.

1914, Tijdschr. Ent. 57: 155 (♀; *Syrphus*). Type-loc: Nongkodjadjar, Java. N. comb.

Genus SCAEVA Fabricius

Scaeva Fabricius, 1805, Syst. Antiatl.: 248. Type-species, *Musca pyrastri* Linnaeus, by designation of Curtis (1834: pl. 509).

Lasiophicus Rondani, 1845, Nuovi Annali Sci. nat. Bologna (2) 2: 459. Type-species, *Musca pyrastri* Linnaeus, by original designation.

Catabomba Osten Sacken, 1877, Bull. U.S. geol. geogr. Surv. Territ. 3: 326. Type-

species, *Musca pyrastri* Linnaeus, by monotypy.

References: Scott 1939; Brauns 1953, 1954 a,b; Dixon 1960 (biol., imm. stages).
latimaculata (Brunetti). — India [Punjab, Rajasthan, Uttar Pradesh], W Pakistan, 1923, Fauna Br. India, Dipt. 3: 68 (♂ ♀; *Lasiopticus*). Type-locs: Allahabad, Peshawar, Ferozepore, and Abu, India.

opimus (Walker). — India.

1852, Insecta Saundersiana 1: 232 (♂; *Syrphus*). Type-loc: East Indies. N. comb.

selenitica (Meigen). — China, India [Assam], Indochina; Europe.

1822, Syst. Beschr. europ. zweifl. Insekts. 3: 304 (sex?; *Syrphus*). Type-loc: Europe.

lunatus Wiedemann, 1830, Aussereurop. zweifl. Insekts. 2: 121 (♀; *Syrphus*). Type-loc: China.

annamites Bigot, 1885, Annls Soc. ent. Fr. (6) 5: 250 (♂; *Lasiophthicus*). Type-loc: Cochinchine.

Genus SPHAEROPHORIA Lepeletier and Serville

Syrphus, subg. *Sphaerophoria* Lepeletier and Serville, in Latreille, et al., 1828, Encycl. méth. (Ins.) 10 (2): 513. Type-species, *Musca scripta* Linnaeus, by designation of Rondani (1844: 458).

Reference: Knutson 1972 (rev., biol.).

assamensis Joseph. — India [Assam].

1970, Eos Madr. 45: 168 (♂ ♀). Type-loc: Dunn Bridge (1524 m), Kameng Frontier Division, Assam, India.

bengalensis Macquart. — India [Himachal Pradesh, Punjab, Uttar Pradesh, W Bengal], Nepal, W. Pakistan [Baluchistan]; Iran.

1842, Dipt. exot. 2(2): 104 (164) (♂). Type-loc: Bengal, India.

flavoabdominalis Brunetti, 1915, Rec. Indian Mus. 11: 214 (♂ ♀; as form). Type-locs: Baluchistan, Bushire, Katmandu (Nepal), Dharampur (Simla Hills), Agra, Ferozepore.

cylindrica (Say). — Not Oriental; Japan, Nearctic.

formosana (Matsumura). — Formosa.

1916, Ent. Mag. Kyoto 2: 23 (♂; *Melithreptus*). Type-loc: Heirinbi, Formosa.

indiana Bigot. — ?Ceylon, China, India [Himachal Pradesh, W Bengal]; Korea, eastern USSR.

1884, Annls Soc. ent. Fr. (6) 4: 99 (♂). Type-loc: "Indes."

nigritarsis Brunetti, 1915, Rec. Indian Mus. 11: 216 (♂ ♀; as form). Type-locs: Simla, and Kurseong, India.

macrogaster (Thomson). — Ceylon, China, India [Madras], Nepal; Australia, Japan, Korea, Manchuria, New Caledonia, New Guinea, eastern USSR.

1869, in K. svenska fregatten Eugenies resa, Zool., Dipt.: 501 (♂ ♀; *Syrphus*). Type-loc: Sydney, Australia.

koreana Bankowska, 1964, Annls zool. Warsz. 22: 339 (♂ ♀). Type-loc: Dephuan Kujang, Korea.

poonaensis Joseph. — India [Maharashtra].

1967, Bull. Ent. Loyola Coll. 8(2): 1 (♀). Type-loc: Botanical Garden, nr River Bhima, Poona, India.

scripta (Linnaeus). — India [Kashmir], Nepal; Palaeartic.

1758, Syst. Nat. Ed. 10: 594 (sex?; *Musca*). Type-loc: Uppsala, Sweden.
brunettii Joseph, 1968, Orient. Insects 1: 243 (♂). Type-loc: Srinagar (1800 m), Kashmir, India.

viridaenea Brunetti. — India.

1915, Rec. Indian Mus. 11: 216 (♂ ♀; as form). Type-loc: Simla (2134 m), India.

vockerothi Joseph. — Formosa, India [Assam].

1970, Eos Madr. 45: 165 (♂ ♀). Type-loc: Kimin (198 m), Subansiri Frontier Division, Assam, India.

Genus SYRPHUS Fabricius

Syrphus Fabricius, 1775, Syst. Ent.: 762. Type-species, *Musca ribesii* Linneaus, by designation of Rondani (1844: 459).

References: Bhatia 1939; Dixon 1960; Dušek and Láska 1964; Scott 1939 (biol., imm. stages).

ribesii (Linneaus). — Not Oriental; North and Central America, Palaeartic.

torvus Osten Sacken. — Formosa, India [Himachal Pradesh], Nepal, Thailand; Holarctic.

1875, Proc. Boston Soc. nat. Hist. 18: 139 (♂ ♀). Type-locs: Canada; Colorado, N.H., Mass., R.I., USA.

vitripennis Meigen. — Formosa; Europe.

1822, Syst. Beschr. zweifl. Insekts. 3: 308 (sex?). Type-loc: Europe.

Unplaced species of Syrphus Fabricius

aeneifrons Brunetti. — India [Assam].

1913, Rec. Indian Mus. 8: 159 (♂). Type-loc: between Kalek and Misshing, NE Frontier, India.

ambiguus Shiraki. — Formosa.

1930, Mem. Fac. Agric. Taihoku imp. Univ. 1: 376 (♂). Type-loc: Rantaisan, Formosa.

angustatus Sack. — Philippines [Luzon].

1926, Philipp. J. Sci. 29: 568 (♂). Type-loc: Luzon, Philippines.

bazini Brunetti. — India [Uttar Pradesh].

1925, Rec. Indian Mus. 27: 75 (n. name for *rufiventris* Brunetti).

rufiventris Brunetti, 1915, Rec. Indian Mus. 11: 202 (♂; *Pipizella*; preocc. Macquart 1849). Type-loc: Kousanie, Kumaon District, India.

duplex Walker. — Malaya.

1857, J. Linn. Soc. Lond. 1: 18 (♂ ♀). Type-loc: Singapore, Malaya.

fulvifacies Brunetti. — China [Yunnan], India [Assam], Java, Laos, Nepal, Philippines [Luzon].

1913, Rec. Indian Mus. 8: 161 (♀). Type-loc: Rotung, NE Frontier, Assam, India.

himalayanus Nayar. — India.

1968, Agra Univ. J. Res. 16: 121 (♂). Type-loc: nr Post Office, Monalsu Gorge, NW Himalaya, India.

isaaci Bhatia. — India [Assam, Bihar, Uttar Pradesh], Nepal.

1933, Indian J. agric. Sci. 2: 566 (♂ ♀). Type-loc: unknown.

issikii Shiraki. — Formosa.

1930, Mem. Fac. Agric. Taihoku imp. Univ. 1: 378 (♂ ♀). Type-locs: Kusukusu, Fuhosho, and Shu-Shu, Formosa.

minimus Shiraki. — Formosa; Japan.

1930, Mem. Fac. Agric. Taihoku imp. Univ. 1: 358 (♂ ♀). Type-locs: Noko, Formosa; Hokkaido, Sapporo, Japan.

neglectus Wiedemann. — Borneo.

1830, Aussereurop. zweifl. Insekten. 2: 134 (♂ ♀). Type-loc: "Vaterland?" (Borneo), Rondani (1875: 423).

remotus Brunetti. — India.

1923, Fauna Br. India, Dipt. 3: 78 (♂). Type-loc: Kufri, Simla Hills, India.

sonami Shiraki. — Formosa.

1930, Mem. Fac. Agric. Taihoku imp. Univ. 1: 389 (♂ ♀). Type-locs: Arisan, Toroen, Asaki, Kanko, and Karenko, Formosa.

ssp. *arisanicus* Shiraki. — Formosa.

1930, Mem. Fac. Agric. Taihoku imp. Univ. 1: 391 (sex?). Type-loc: Arisan, Formosa.

transversus Brunetti. — India [Assam].

1913, Rec. Indian Mus. 8: 160 (♀). Type-loc: Sadiya, India.

Nomen nudum

instabilis Brunetti (MS), Bhatia and Shaffi, 1933, Indian J. agric. Sci. 2: 566.

Genus XANTHOGRAMMA Schiner

Xanthogramma Schiner, 1860, Wien. ent. Mschr. 4: 215. Type-species, *Syrphus ornatus* Meigen, 1822, by designation of Williston (1887: 91), = *peditsequeum* (Harris).

Olibrosyrphus Mik, 1897, Wien. ent. Ztg 16: 66. Type-species, *Syrphus laetus* Fabricius, by monotypy.

Schiner proposed *Xanthogramma* without a binomen and without a type-species; therefore the gender may be considered feminine by reason of Schiner (1861: 318).

arisanica Shiraki. — Formosa.

1930, Mem. Fac. Agric. Taihoku imp. Univ. 1: 405 (♂ ♀). Type-locs: Arisan, Shishito, Kosempo, and Fuhosho, Formosa.

calceata Sack. — Philippines [Negros].

1926, Philipp. J. Sci. 29: 570 (♀). Type-loc: Canlaon Volcano, Negros Occidental, Philippines.

fasciata Shiraki. — Formosa.

1930, Mem. Fac. Agric. Taihoku imp. Univ. 1: 410 (♂ ♀). Type-locs: Rakuraku, and Kosempo, Formosa.

fumipennis Matsumura. — Formosa.

1916, Ent. Mag. Kyoto 2(1): 29 (♂). Type-loc: Horisha, Formosa.

indica Nayar. — India [Himachal Pradesh].

1968, Agra Univ. J. Res. 16: 129 (♀). Type-loc: Kalatop, nr Dalhousie, India.

pruthii Deoras. — India [Uttar Pradesh].

1942, Indian J. Ent. 4: 217 (♂ ♀). Type-loc: Delhi, India.

Tribe BACCHINI

Genus BACCHA Fabricius

Baccha Fabricius, 1805, Syst. Antlat.: 199. Type-species, *Syrphus elongatus* Fabricius, by designation of Curtis (1839: pl. 737).

References: Curran 1947; de Meijere 1908 (key); Bhatia and Shaffi 1933 (biol., imm. stages).

Subgenus ALLOBACCHA Curran

Baccha, subg. *Allobaccha* Curran, 1928, J. fed. Malay St. Mus. 14: 251. Type-species, *Baccha rubella* van der Wulp, by original designation.

amphitheo Walker. — Ceylon, India [Assam], Formosa south to Flores.

1849, List Dipt. Br. Mus. 3: 549 (sex?). Type-loc: "E. Ind./ Mulmein." *pedicellata* Doleschall, 1856, Natuurk. Tijdschr. Ned.-Indië 10: 411 (sex?). Type-loc: Java.

bicincta de Meijere, 1910, Tijdschr. Ent. 53: 104 (♂ ♀). Type-locs: Batavia (Djakarta), Tandjong Priok, Bekassi, Java and Krakatau.

flavopunctata Brunetti, 1913, Rec. Indian Mus. 8: 165 (♀). Type-loc: Dibrugarh, Assam, India.

fulvicostalis Matsumura, 1916, Thousand Insects Japan 2: 226 (♀). Type-loc: Formosa.

apicalis Loew. — Ceylon, Formosa, India [Uttar Pradesh], Java, Malaya, Philippines; Bonin Islands, Japan.

1858, Wien. ent. Mschr. 2: 106 (♀). Type-loc: Japan.

pulchrifrons Austen, 1893, Proc. zool. Soc. Lond. 1893: 139 (♂ ♀). Type-loc: Hot Wells, Trincomali, Ceylon.

apicenotata Brunetti, 1915, Rec. Indian Mus. 11: 221 (♀). Type-loc: Bhowali (5700 ft), W Himalayas (Kumaon), India.

bicolor Austen. — Sula; Moluccas (Mysol), New Guinea.

1893, Proc. zool. Soc. Lond. 1893: 137 (♂). Type-loc: Mysol Island, Moluccas.

bimaculata Keiser. — Sumbawa.

1952, Verh. naturf. Ges. Basel 63: 161 (♀). Type-loc: Langgai, central Sumbawa.

chalybaea de Meijere. — Java, Krakatau.

1910, Tijdschr. Ent. 53: 105 (♂ ♀). Type-locs: Pasuruan, Java; and Krakatau.

chalybescens Curran. — Malaya.

1931, J. fed. Malay St. Mus. 16: 326 (♀). Type-loc: Bukit Kutu (3500 ft), Selangor, Malaya.

circumcincta de Meijere. — Java.

1908, Tijdschr. Ent. 51: 320 (♀). Type-loc: Buitenzorg (Bogor), Java.

- cochleariformis* Sack. — Philippines [Mindanao].
1926, Philipp. J. Sci. 29: 573 (♀). Type-loc: Surigao, Mindanao, Philippines.
- dispar* Walker. — Celebes, India, Malaya.
1860, J. Linn. Soc. Lond. 4: 121 (♂). Type-loc: Celebes.
- fallax* Austen. — Borneo, Ceylon, Formosa, India, Malaya; Japan.
1893, Proc. zool. Soc. Lond. 1893: 142 (♂). Type-loc: Haycock Hill, nr Galle, Ceylon.
- gigas* Curran. — Borneo.
1931, J. fed. Malay St. Mus. 16: 356 (♀). Type-loc: Mt Kinabalu, Marei Parei (5000 ft), N Borneo.
- keiseri* van der Goot. — Sumbawa.
1964, Beaufortia 10: 219 (n. name for *violacea* Keiser).
violacea Keiser, 1952, Verh. naturf. Ges. Basel 63: 163 (♂; preocc. Hull 1943). Type-loc: Langgaliru, Sumbawa.
- luteolimbata* de Meijere. — Sumatra.
1924, Tijdschr. Ent. 67 (Suppl): 22 (♂ ♀). Type-loc: "Aur Kumanis" (Sumatra).
- macgregori* Curran. — Philippines [Luzon, Calicoan].
1929, Ann. ent. Soc. Am. 22: 492 (♂ ♀). Type-loc: Manila, Philippines.
- meijerei* Kertész. — Formosa, Java, Krakatau, Philippines [Mindanao], Simeuluë Island, Sumbawa; Moluccas (Buru).
1913, Annls hist.-nat. Mus. natn. hung. 11: 279 (♂ ♀). Type-locs: Tandjong Priok and Semarang, Java; Krakatau.
pedicellata, of de Meijere, not Doleschall.
- nigrapex* Curran. — Malaya.
1931, J. fed. Malay St. Mus. 16: 324 (♂). Type-loc: Bukit Kutu (3500 ft), Selangor, Malaya.
- nigriceps* de Meijere. — Sumatra.
1924, Tijdschr. Ent. 67 (Suppl): 23 (♂). Type-loc: Kadjai (Ophir District), Sumatra.
- nigricosta* Brunetti. — India [Uttar Pradesh], W Pakistan.
1907, Rec. Indian Mus. 1: pl. xi, fig. 5; 1908, Rec. Indian Mus. 2: 50 (♂). Type-loc: Bhim Tal, Kumaon, India.
- nigricoxa* Curran. — Malaya.
1928, J. fed. Malay St. Mus. 14: 249 (♂). Type-loc: Rhododendron Hill, Cameron Highlands, Pahang, Malaya.
- nubilipennis* Austen. — Ceylon, Formosa, India [Assam], Nepal; Japan.
1893, Proc. zool. Soc. Lond. 1893: 136 (♂ ♀). Type-loc: Kandy, Ceylon.
- purpuricola* Walker. — Philippines [Luzon, Negros]; Moluccas (Key, Mysol), New Guinea.
1859, J. Linn. Soc. Lond. 3: 129 (♀). Type-loc: Key Island.
- refulgens* Austen. — Java, Sumbawa; Moluccas (Buru).
1893, Proc. zool. Soc. Lond. 1893: 138 (♀). Type-loc: Buru Island, Moluccas.
- sapphirina* Wiedemann. — Ceylon, India [Madras, Maharashtra, Rajasthan], Formosa; New Guinea, Aden, Africa.
1830, Aussereurop. zweifl. Insekt. 2: 96 (♀). Type-loc: East Indies.

- signata* Sack. — Philippines [Mindanao].
1926, Philipp. J. Sci. 29: 575 (♂). Type-loc: Kolambugan, Lanao, Mindanao, Philippines.
- sulica* Austen. — Sula; Moluccas (Mysol).
1893, Proc. zool. Soc. Lond. 1893: 144 (♀). Type-locs: Sula; and Mysol, Moluccas.
- sumbana* Keiser. — Sumbawa.
1952, Verh. naturf. Ges. Basel 63: 163 (♂). Type-loc: Mangrovewald, Melolo, Sumbawa.
- tinctiventris* de Meijere. — Sumatra.
1924, Tijdschr. Ent. 67 (Suppl): 21 (♀). Type-loc: "Aur Kumanis" (Sumatra).
- triangulifera* Austen. — Ceylon, India [Sikkim, W Bengal], Nepal.
1893, Proc. zool. Soc. Lond. 1893: 138 (♂ ♀). Type-loc: Huldamulla, Ceylon.
- elegans* Brunetti, 1915, Rec. Indian Mus. 11: 220 (♂). Type-loc: Sukna, base of Darjeeling Hills, India.
- tripartita* Walker. — Celebes, Sula; Moluccas.
1861, J. Linn. Soc. Lond. 5: 285 (♂). Type-loc: Batjan, Moluccas.
- umbrifera* de Meijere. — Sumatra.
1924, Tijdschr. Ent. 67 (Suppl): 23 (♂). Type-loc: Kadjai (Ophir District), Sumatra.
- umbrosa* Brunetti. — India [Rajasthan].
1923, Fauna Br. India, Dipt. 3: 119 (♂). Type-loc: Abu, Rajputana, India.
- varipes* Curran. — Malaya.
1928, J. fed. Malay St. Mus. 14: 252 (♂). Type-loc: Seat Point (5460 ft), Gunong Tahan, Pahang, Malaya.
- wegneri* Keiser. — Sumbawa.
1952, Verh. naturf. Ges. Basel 63: 164 (♀). Type-loc: Mangrovewald, Melolo, Sumbawa.
- ### Subgenus BACCHA Fabricius
- maculata* Walker. — Borneo, Formosa, India [Assam, Himachal Pradesh, Uttar Pradesh, W Bengal], Java, Malaya, Nepal, Philippines [Luzon], Sumatra; Japan, Korea.
1852, Insecta Saundersiana 1: 223 (♂). Type-loc: East Indies.
- tinctipennis* Brunetti, 1907, Rec. Indian Mus. 1: pl. xi, fig. 6; 1908, Rec. Indian Mus. 2: 51 (♀). Type-loc: Bhim Tal, India.
- austeni* de Meijere, 1908, Tijdschr. Ent. 51: 325 (♂ ♀). Type-loc: Gunung Salak nr Buitenzorg (Bogor), Java.
- tenera* de Meijere, 1910, Tijdschr. Ent. 53: 103 (♂ ♀). Type-loc: Tankuban Prahu, Java.
- eronis* Curran, 1928, J. fed. Malay St. Mus. 14: 248 (♂). Type-loc: Lubok Tamang (3500 ft), Pahang, Malaya.
- virtuosa* Curran. — Malaya.
1928, J. fed. Malay St. Mus. 14: 246 (♂). Type-loc: Rhododendron Hill (5200 ft), Cameron Highlands, Pahang, Malaya.

Unplaced species of *Baccha* Fabricius

- brunnea* Hull. — Java.
1949, Ann. Mag. nat. Hist. (12) 2: 733 (♂). Type-loc: Sockaboemi, Java.
- gratiosa* Bigot. — Borneo.
1883, Annls Soc. ent. Fr. (6) 3: 335 (♀). Type-loc: Sarawak, Borneo.
- loriae* de Meijere. — Philippines [Luzon]; New Guinea.
1908, Tijdschr. Ent. 51: 324 (♀). Type-loc: Paumomu, New Guinea.
- mundula* van der Wulp. — Java; New Guinea.
1898, Természetr. Füz. 21: 423 (♀). Type-loc: New Guinea.
- plumbicincta* Brunetti. — India [Assam].
1915, Rec. Indian Mus. 11: 222 (♀). Type-loc: Cherrapunji, Assam, India.
- porphyra* Curran. — Borneo, Malaya.
1928, J. fed. Malay St. Mus. 14: 254 (♂ ♀). Type-loc: Wray's Camp (3300 ft), Pahang, Malaya.
- sauteri* Kertész. — Formosa.
1913, Annls hist.-nat. Mus. natn. hung. 11: 275 (♂ ♀). Type-locs: Takao, Janano-Taiko, Kosempo, Tainan, Koshun, Fuhosho, and Sokotsu, Formosa.
- tribeni* Nayar. — India [Himachal Pradesh].
1968, Agra Univ. J. Res. 16: 128 (♂). Type-loc: 67 km on Hindustan-Tibet Rd, from Narkanda, Simla, India.
- vespaiformis* Doleschall. — Celebes; Moluccas (Amboina).
1857, Natuurk. Tijdschr. Ned.-Indië 14: 411 (sex?). Type-loc: Amboina, Moluccas.
- vivida* Hull. — Timor.
1944, Ann. Mag. nat. Hist. (11) 11: 50 (♀). Type-loc: Timor, Dutch East Indies (Indonesia).

Genus RHINOBACCHA de Meijere

- Rhinobaccha* de Meijere, 1908, Tijdschr. Ent. 51: 315. Type-species, *gracilis* de Meijere, by monotypy.
- gracilis* de Meijere. — Ceylon.
1908, Tijdschr. Ent. 51: 315 (♂ ♀). Type-loc: Pattipola, Ceylon.

Tribe MELANOSTOMATINI

Genus MELANOSTOMA Schiner

- Melanostoma* Schiner, 1860, Wien. ent. Mschr. 4: 213. Type-species, *Musca mellina* Linneaus, by monotypy.
- Reference: Dušek and Láska 1959 (biol., imm. stages).
- abdominale* Shiraki. — Formosa.
1930, Mem. Fac. Agric. Taihoku imp. Univ. 1: 330 (♂ ♀). Type-locs: Arisan and Musha, Formosa.

algens Curran. — Borneo.

1931, J. fed. Malay St. Mus. 16: 359 (♂ ♀). Type-loc: Mt Kinabalu, Pakka (10,000 ft), N Borneo.

ambiguum (Fallén). — India [Himachal Pradesh].

1817, Syrphici Sveciae: 47 (♀; *Syrphus*). Type-loc: Europe.

atrum Sack. — Lombok.

1932, Zool. Anz. 100: 227 (♂ ♀). Type-loc: Segare Anak, Lombok, Indonesia.

gedehense de Meijere. — Java.

1914, Tijdschr. Ent. 57: 163 (♂ ♀; as *quadrinotatum* var.). Type-loc: Gunung Gede, Java.

meijerei van der Goot. — Java.

1964, Beaufortia 10: 219 (n. name for *quadrinotatum* de Meijere). *quadrinotatum* de Meijere, 1914, Tijdschr. Ent. 57: 163 (♂ ♀; preocc. Bigot 1884). Type-loc: Nongkodjadjar, Java.

normale Curran. — Borneo.

1931, J. fed. Malay St. Mus. 16: 361 (♂ ♀). Type-loc: Mt Kinabalu, Marei Parei, N Borneo.

orientale (Wiedemann). — throughout Oriental region.

1824, Analecta Ent.: 36 (♀; *Syrphus*). Type-loc: "Ind. Or."

pedius (Walker). — ?Java.

1852, Insecta Saundersiana 1: 234 (♂; *Syrphus*). Type-loc: East Indies.
cothonea Walker, 1852, Insecta Saundersiana 1: 235 (♀; *Syrphus*). Type-loc: East Indies.

quadrifasciatum Curran. — Malaya.

1928, J. fed. Malay St. Mus. 14: 260 (♀). Type-loc: Jor Camp, (1800 ft), Batang Padang, Perak, Malaya.

scalare (Fabricius). — throughout Oriental region; New Guinea, Palaearctic.

1794, Ent. Syst. 4: 308 (sex?; *Syrphus*). Type-loc: "Kiliae."
ceylonense de Meijere, 1911, Tijdschr. Ent. 54: 348 (♂ ♀). Type-loc: Pattipola (2000 m), Ceylon.

univittatum (Wiedemann). — throughout SE Asia; ?Australia, ?Fiji, Japan, ?New Caledonia, ?Samoa.

1824, Analecta Ent.: 36 (sex?; *Syrphus*). Type-loc: "Ind. Or."

gathifer Walker, 1857, J. Linn. Soc. Lond. 1: 125 (♀; *Syrphus*). Type-loc: Sarawak, Borneo.

planifacies Macquart, 1848, Dipt. exot. Suppl. 3: 43 (203) (♀; *Syrphus*). Type-loc: Java.

Genus PLATYCHEIRUS Lepeletier and Serville

Platycheirus Lepeletier and Serville, in Latreille, et al., 1828, Encycl. Méth. (Ins.) 10(2): 513. Type-species, *Syrphus scutatus* Meigen, by designation of Westwood (1840: 137).

albimanus (Fabricius). — ?India [Himachal Pradesh, W Bengal], ?Nepal, Philippines; Holarctic.

1781, Spec. Insect. 2: 434 (sex?; *Syrphus*). Type-loc: "Angliae."

formosanus Shiraki. — Formosa.

1930, Mem. Fac. Agric. Taihoku imp. Univ. 1: 325 (♂ ♀). Type-locs: Arisan, Royeichi, Shukoran, Taiheisan, Nitakayama, Rantaisan, Asahi,

Daisukutsu, Musha, Funkiko, Hassensan, Kanko.

himalayensis Brunetti. — India [Uttar Pradesh]. Nepal.

1915, Rec. Indian Mus. 11: 209 (♂; as *manicatus* var.). Type-loc: Garhwal, W Himalayas. N. status.

Genus XANTHANDRUS Verrall

Xanthandrus Verrall, 1901, British Flies 8: 316. Type-species, *Musca comtus* Harris, by designation of Coquillett (1910: 620).

Reference: Dušek and Láska 1967 (biol., imm. stages).

callidus Curran. — Malaya, Thailand.

1928, J. fed. Malay St. Mus. 14: 261 (♂). Type-locs: Khao Luang (5800 ft), Nakhon Si Thammarat, Thailand; Rhododendron Hill, Cameron Highlands, Pahang (5500 ft) and Gunong Tahan, Pahang, Malaya.

ceylonicus Keiser. — Ceylon.

1958, Revue suisse Zool. 65: 205 (♂). Type-loc: Pidratalagala, Ceylon. *comtus* (Harris). — Formosa; Europe, Japan.

?1780, Expos. engl. Insects: 108 (sex?; *Musca*). Type-loc: England.

indicus Curran. — India [Madhya Pradesh].

1933, Stylops 2: 46 (♂ ♀). Type-loc: Rahatgaon, Hoshangabad, India.

orientalis Sack. — Philippines [Negros].

1926, Philipp. J. Sci. 29: 566 (♂ ♀). Type-loc: Canlaon, Negros Occidental, Philippines.

talamaui (de Meijere). — Malaya, Sumatra.

1924, Tijdschr. Ent. 67 (Suppl. 15): 21 (♀; *Melanostoma*). Type-loc: Gunung Talamau, Sumatra.

brevicornis Curran, 1928, J. fed. Malay St. Mus. 14: 263 (♂). Type-loc: Pahang, Cameron Highlands, Rhododendron Hill, Malaya.

Tribe CHRYSOTOXINI

Genus CHRYSOTOXUM Meigen

Antiope Meigen, 1800, Nouv. Class. Mouches: 32. Type-species, *Musca bicincta* Linnaeus, by designation of Coquillett (1910: 508). Suppressed by I.C.Z.N. 1963, Opinion 678: 339.

Chrysotoxum Meigen, 1803, Magazin Insektenkde 2: 275. Type-species, *Musca bicincta* Linnaeus, by designation of Latreille (1810: 443).

Chrysotoxum, subg. *Primochrysotoxum* Shannon, 1926, Proc. U.S. natn. Mus. 69 (11): 2. Type-species, *ypsilone* Williston, by original designation.

antiquum Walker. — India [Himachal Pradesh].

1852, Insecta Saundersiana 1: 218 (♂). Type-loc: East Indies.

atratum de Meijere. — Sumatra.

1921, Tijdschr. Ent. 64: lii (♂). Type-loc: Sumatra.

baphyrus Walker. — Ceylon, India [Uttar Pradesh], Indochina, ?Java, Laos, Nepal.

1849, List Dipt. Br. Mus. 3: 542 (♂). Type-loc: north Bengal, India.

indicum Walker, 1852, Insecta Saundersiana 1: 218 (♂). Type-loc: East Indies.

sexfasciatum Brunetti, 1907, Rec. Indian Mus. 1: pl. xiii, fig. 9; 1908: 89 (♀).

Type-loc: Bijnor District, India.

citronellum Brunetti, 1908, Rec. Indian Mus. 2: 90 (♂). Type-loc: Kandy, Ceylon.

mundulum Hervé-Bazin, 1923, Bull. Soc. ent. Fr. 1923: 27 (♂). Type-loc: Cochinchina.

convexum Brunetti. — India [Uttar Pradesh].

1915, Rec. Indian Mus. 1: 249 (♂). Type-loc: Andarban, Garhwal District, W Himalayas, India.

festivum (Linnaeus). — India [Bihar]; Palaearctic.

1758, Syst. Nat. Ed. 10: 593 (sex?; *Musca*). Type-loc: Europe.

formosanum Shiraki. — Formosa.

1930, Mem. Fac. Agric. Taihoku imp. Univ. 1: 27 (♂ ♀). Type-locs: Arisan, and Paiwan, Formosa.

fuscomarginatum Brunetti. — W Pakistan.

1923, Fauna Br. India, Dipt. 3: 300 (♀). Type-loc: Chitral (Pakistan).

ladakense Shannon. — India [Kashmir].

1926, Proc. U.S. natn. Mus. 69 (11): 13 (♀). Type-loc: Rupshu Ledak (16,000 ft), Kashmir, India.

quadrifasciatum Brunetti. — India [Assam], Laos.

1923, Fauna Br. India, Dipt. 3: 300 (♀). Type-loc: Khasi Hills, Assam, India.

rotundatum Hervé-Bazin, 1923, Bull. Soc. ent. Fr. 1923: 27 (♀). Type-loc: Laos.

testaceum Sack. — Formosa; Japan.

1913, Ent. Mitt. 2: 9 (♂ ♀). Type-locs: Yama, and Tappani, Formosa.

violaceum Brunetti. — India [W Bengal].

1923, Fauna Br. India, Dipt. 3: 302 (♂ ♀). Type-loc: Sureil, Darjeeling District, India.

Tribe PARAGINI

Genus PARAGUS Latreille

Paragus Latreille, 1804, Nouv. Dict. Hist. nat. 24 (Tab. Méth. Ins.): 94. Type-species, *Syphus bicolor* Fabricius, by monotypy.

References: Stuckenberg 1954 (key); Bhatia and Shaffi 1933; Heiss 1938 (biol., imm. stages).

Subgenus PARAGUS Latreille

auritus Stuckenberg. — Ceylon, India [W Bengal], Nepal; Kenya.

1954, Trans. R. ent. Soc. Lond. 105: 418 (♂ ♀). Type-loc: Kandy, Ceylon.

crenulatus Thomson. — entire Oriental region; Australia.

1869, in K. svenska fregatten Eugenies resa, Zool., Dipt.: 503 (♂). Type-loc: China.

serratus (Fabricius). — India [Madhya Pradesh, Madras, Maharashtra, Mysore, Punjab].

1805, Syst. Antliat.: 186 (sex?; *Mulio*). Type-loc: Tranquebar, India.

yerburiensis Stuckenberg. — Ceylon, India [Madhya Pradesh], Nepal.
1954, Trans. R. ent. Soc. Lond. 105: 415 (♂ ♀). Type-loc: Velverry,
Ceylon.

Subgenus PANDASYOPHTHALMUS Stuckenberg

Pandasyophthalmus Stuckenberg, 1954, Revue Zool. Bot. afr. 49: 100. Type-species, *longiventris* Loew, by original designation.

atratus de Meijere. — India [Uttar Pradesh], Java, Malaya, Sumatra, Sumbawa, Timor; New Guinea.

1906, Nova Guinea 1: 85 (♂). Type-loc: Manokwari, New Guinea.

hokusankoensis Shiraki. — Formosa.

1930, Mem. Fac. Agric. Taihoku imp. Univ. 1: 249 (♂ ♀). Type-loc: Hokusanko, Formosa.

keiseri van der Goot. — Java, Sumbawa.

1964, Beaufortia 10: 219 (n. name for *ruficaudatus* Keiser).

ruficaudatus Keiser, 1952, Verh. naturf. Ges. Basel 63: 154 (♂; preocc. Bigot 1884). Type-loc: Lokojengo, Sumbawa.

politus Wiedemann. — China, Philippines.

1830, Aussereurop. zweifl. Insekt. 2: 89 (sex?). Type-loc: China.

rufocincta (Brunetti). — Burma, India.

1908, Rec. Indian Mus. 2: 53 (♂; *Pipizella*). Type-loc: Rangoon, Burma.

tibialis (Fallén). — widespread Oriental; cosmopolitan.

1817, Syrphici Sveciae: 60 (♂ ♀; *Pipiza*). Type-loc: "Vestergothia," Sweden.

indicus Brunetti, 1908, Rec. Indian Mus. 2: 52 (♂ ♀; *Pipizella*). Type-loc: Matiana, Simla, India.

ssp. *rufiventris* Brunetti. — Ceylon, India [Assam], Nepal, W Pakistan.

1913, Rec. Indian Mus. 8: 157 (♂; as sp.). Type-loc: Mangaldai, Assam-Bhutan Frontier, India.

Unplaced species of *Paragus* Latreille

latiusculus Walker. — Celebes.

1861, J. Linn. Soc. Lond. 5: 266 (♀). Type-loc: Tond, Celebes.

Subfamily MILESIINAE

Tribe PIPIZINI

Genus PIPIZA Fallén

Pipiza Fallén, 1810. Sp. Ent. nov. Dipt.: 11. Type-species, *Musca noctiluca* Linnaeus, by designation of Curtis (1837: pl. 669).

signata Meigen. — Formosa; Palaearctic.

1822, Syst. Beschr. europ. zweifl. Insekt. 3: 246 (♂ ♀). Type-loc: Europe.

Genus TRIGLYPHUS Loew

Triglyphus Loew, 1840, Isis (Oken's) 1840: 512. Type-species, *prinii* Loew, by monotypy.

cyanus (Brunetti). — India [Sikkim].

1915, Rec. Indian Mus. 11: 202 (♀; *Psilota*). Type-loc: Gangtok, Sikkim, India.

formosanus Shiraki. — Formosa.

1930, Mem. Fac. Agric. Taihoku imp. Univ. 1: 251 (♀). Type-loc: Mu-sha, Formosa.

Tribe CHEILOSIINI

Genus CHEILOSIA Meigen

Cheilosia Meigen, 1822, Syst. Beschr. europ. zweifl. Insekt. 3: 296. Type-species, *Eristalis scutellatus* Fallén, by designation of Rondani (1856: 51).

Chilosia, emendation. (see Stone, et al. 1965: 583).

acuticornis Curran. — Malaya.

1928, J. fed. Malay St. Mus. 14: 267 (♂ ♀). Type-locs: Ginting Sempak, Pahang border, Selangor and Lubok Tamang, Pahang, Malaya.

apicalis Brunetti. — India [Assam].

1913, Rec. Indian Mus. 8: 158 (♀). Type-loc: Rotung, NE Frontier, Assam, India.

atterima Sack. — Formosa; Japan.

1927, Stettin. ent. Ztg 88: 305 (♂). Type-loc: Toa Tsu Kutsu, Formosa.

bombylioides Hull. — Burma.

1949, Ann. Mag. nat. Hist. (12) 2: 739 (♀). Type-loc: Adung Valley, N Burma.

formosana Shiraki. — Formosa.

1930, Mem. Fac. Agric. Taihoku imp. Univ. 1: 309 (♂ ♀). Type-loc: Rikiriki, Funkiko, Niitakagun, Formosa.

grossa (Fallén). — India [Uttar Pradesh].

1817, Syrphici Sveciae: 53 (♀; *Eristalis*). Type-loc: Europe.

himalayensis (Brunetti). — India [Uttar Pradesh].

1915, Rec. Indian Mus. 11: 217 (♂; *Eriozona*). Type-locs: Kumaon District, W Himalayas and Onari, Garhwal District, India.

hirticincta Brunetti. — India [W Bengal].

1915, Rec. Indian Mus. 11: 204 (♂). Type-loc: Darjeeling District, India.

kalatopensis Nayar. — India [Punjab].

1968, Agra Univ. J. Res. 16: 129 (♀). Type-loc: Kalatop-Lakkarmandi Bridal path, India.

leptorrhyncha Shiraki. — Formosa.

1930, Mem. Fac. Agric. Taihoku imp. Univ. 1: 269 (♂ ♀). Type-locs: Shukoran, Kanko, Daisuikutsu, and Roeichi, Formosa.

makiana Shiraki. — Formosa.

1930, Mem. Fac. Agric. Taihoku imp. Univ. 1: 272 (♀). Type-locs: Karenko, and Horisha, Formosa.

- nigroaenea* Brunetti. — India, Nepal.
1915, Rec. Indian Mus. 11: 204 (♂ ♀). Type-loc: Simla District, India.
- niitakana* Shiraki. — Formosa.
1930, Mem. Fac. Agric. Taihoku imp. Univ. 1: 275 (♀). Type-loc: Niita-kayama (4200 m), Formosa.
- ochreipila* Shiraki. — Formosa.
1930, Mem. Fac. Agric. Taihoku imp. Univ. 1: 283 (♂ ♀). Type-locs: Shukoran, Taiheisan, and Rocichi, Formosa.
- pilipes* (Bigot). — India.
1883, Annls Soc. ent. Fr. (6) 3: 551 (♀; *Cartosyrphus*). Type-loc: India.
- plumbiventris* Brunetti. — India.
1915, Rec. Indian Mus. 11: 205 (♀). Type-loc: Simla, India.
- rakurakuensis* Shiraki. — Formosa.
1930, Mem. Fac. Agric. Taihoku imp. Univ. 1: 265 (♂ ♀). Type-loc: Rakuraku, Formosa.
- splendida* Shiraki. — Formosa.
1930, Mem. Fac. Agric. Taihoku imp. Univ. 1: 302 (♂ ♀). Type-locs: Karenko, Asahi, and Musha, Formosa.
- tendens* Curran. — Malaya.
1928, J. fed. Malay St. Mus. 14: 269 (♂). Type-loc: Lubok Tamang (3500 ft), Pahang, Malaya.
- yoshinoi* Shiraki. — Formosa.
1930, Mem. Fac. Agric. Taihoku imp. Univ. 1: 267 (♀). Type-loc: Karenko, Formosa.

Genus ENDOIASIMYIA Bigot

- Endoiasimyia* Bigot, 1882, Annls Soc. ent. Fr. (6) 2 (Bull.); cxxxvi. Type-species, *indiana* Bigot, by monotypy.
- Sonanomyia* Shiraki, 1930, Mem. Fac. Agric. Taihoku imp. Univ. 1: 320. Type-species, *formosana* Shiraki, by original designation.
- formosana* (Shiraki). — Formosa.
1930, Mem. Fac. Agric. Taihoku imp. Univ. 1: 321 (♂ ♀; *Sonanomyia*). Type-locs: Asahi, Tenchosan, Arisan, and Shu-Shu, Formosa.
- indiana* Bigot. — India, Nepal.
1882, Annls. Soc. ent. Fr. (6) 2 (Bull.); cxxxvi (♀). Type-loc: India.
- javanensis* (de Meijere). — Java, Sumatra.
1914, Tijdschr. Ent. 57: 150 (♀; *Cheilosia*). Type-locs: Nongkodjadjar, Gunung Ungaran, and Gunung Gede, Java.

Genus FERDINANDEA Rondani

- Ferdinandea* Rondani, 1844, Nuovi Annali Sci. nat. Bologna (2) 2: 196. Type-species, *Conops cuprea* Scopoli, by designation of Rondani (1856: 56).
References: Coe 1964; Hull 1942 (keys).
- formosana* Shiraki. — Formosa.
1930, Mem. Fac. Agric. Taihoku imp. Univ. 1: 103 (♂). Type-loc: Mu-sha, Formosa.

- isabella* Hull. — India [Kashmir].
1942, J. Wash. Acad. Sci. 32: 241 (♀). Type-loc: Gulmarg (8500 ft), Kashmir, India.
- longifacies* Coe. — Nepal.
1964, Bull. Br. Mus. nat. Hist. (Ent.) 15: 266 (♂ ♀). Type-loc: Taplejung District, above Sangu, E Nepal.
- maculipennis* Curran. — Malaya.
1928, J. fed. Malay St. Mus. 14: 271 (♂). Type-loc: Gunong Berumban (6050 ft), Cameron Highlands, Pahang, Malaya.
- montana* Hull. — India [Kashmir].
1942, J. Wash. Acad. Sci. 32: 240 (♂ ♀). Type-loc: Gulmarg, Kashmir (8500 ft), India.
- sumatrensis* de Meijere. — Sumatra.
1924, Tijdschr. Ent. 67 (Suppl.): 24 (♂). Type-loc: Gunung Talamau, Sumatra.

Genus RHINGIA Scopoli

- Rhingia* Scopoli, 1763, Ent. Carniolica: 358. Type-species, *Conops rostrata* Linnaeus, by monotypy.
- angusticincta* Brunetti. — India [W Bengal].
1908, Rec. Indian Mus. 2: 59 (♂ ♀). Type-locs: between Kufi and Simla; Theog, India.
- basalis* de Meijere. — Sumatra.
1924, Tijdschr. Ent. 67: 219 (♂). Type-loc: Gunung Singalang, Sumatra.
- binotata* Brunetti. — Formosa, India [Assam, W Bengal], Nepal.
1908, Rec. Indian Mus. 2: 59 (♂). Type-loc: Darjeeling, India.
ssp. *quadrinotata* Hervé-Bazin. — India.
1914, Insecta 41: 151 (sex?). Type-loc: "Inde."
- chaetosa* Curran. — Malaya.
1928, J. fed. Malay St. 14: 275 (♀). Type-loc: Padang (5500 ft), Gunong Tahan, Pahang, Malaya.
- cincta* de Meijere. — Formosa, India [Assam, W Bengal], Java.
1904, Bijdr. Dierk. 18: 101 (♀). Type-loc: Java.
- formosana* Shiraki. — Formosa.
1930, Mem. Fac. Agric. Taihoku imp. Univ. 1: 431 (♂ ♀). Type-locs: Arisan, Musha, Tainan, Kanko, Taiheisan, Asahi, Dajurin, Rantaisan, and Karenko, Formosa.
- laticincta* Brunetti. — India [Himachal Pradesh, Punjab, Uttar Pradesh, W Bengal].
1907, Rec. Indian Mus. 1: pl. xi, figs. 7 & 8; 1908, Rec. Indian Mus. 2: 58 (♂ ♀). Type-locs: Darjeeling District, and Mussoorie, India.
ssp. *fasciata* Brunetti. — India [Punjab, W Bengal].
1908, Rec. Indian Mus. 2: 58 (♂ ♀). Type-loc: Darjeeling, India.
- robinsoni* Edwards. — Sumatra.
1919, J. fed. Malay St. Mus. 8: 38 (♂). Type-loc: Korinchi Peak, Sumatra.
- semincinea* Brunetti. — India [Uttar Pradesh].
1923, Fauna Br. India, Dipt. 3: 134 (♂). Type-loc: Naini Tal, India.

sexmaculata Brunetti. — India [Assam].

1913, Rec. Indian Mus. 8: 166 (♀). Type-loc: Dibrugarh, Assam, India.
siwalikensis Nayar. — India [Punjab].

1968, Agra Univ. J. Res. 16: 126 (♂). Type-loc: Kalatop, nr Dalhousie, India.

uniformis Curran. — Borneo.

1931, J. fed. Malay St. Mus. 16: 365 (♂ ♀). Type-loc: Mt Kinabalu, Kamborangah (7200 ft), N Borneo.

varipennis Curran. — Borneo.

1931, J. fed. Malay St. Mus. 16: 364 (♂ ♀). Type-loc: Mt Kinabalu, Lumu Lumu (5000–5500 ft), N Borneo.

Tribe CALLICERINI

Genus CALLICERA Panzer

Callicera Panzer, 1809, Fauna Germanica 104: 17. Type-species, *Bibio aenea* Fabricius, by monotypy.

Reference: Coe 1964 (key).

aenea (Fabricius). — Formosa; Europe.

1781, Spec. Insect. 2: 413 (sex?; *Bibio*). Type-loc: Germany.

doleschalli Verrall. — Burma.

1913, Trans. ent. Soc. Lond. 1913: 328 (♂). Type-loc: N Chin Hills (5000 ft), Burma.

nitens Coe. — Nepal.

1964, Bull. Br. Mus. nat. Hist. (Ent.) 15: 287 (♀). Type-loc: above Sangu, Taplejung District, E Nepal.

robusta Coe. — India [Uttar Pradesh].

1964, Bull. Br. Mus. nat. Hist. (Ent.) 15: 286 (♂ ♀). Type-loc: Naini Hills (7000–8000 ft), NE India.

sackeni Verrall. — Burma.

1913, Trans. ent. Soc. Lond. 1913: 331 (♂). Type-loc: N Chin Hills, Burma.

sumatrensis de Meijere. — Borneo, Malaya, Sumatra.

1919, Bijdr. Dierk. 21: 26 (♀). Type-loc: Suban Ajam, Sumatra.

pendleburyi Curran, 1928, J. fed. Malay St. Mus. 14: 273 (♂). Type-locs: Cameron Highlands; Rhododendron Hill (5200 ft); Gunong Berumban (summit) (6050 ft) and (5100 ft), Pahang, Malaya.

Tribe VOLUCELLINI

Genus GRAPTONYZA Wiedemann

Graptomyza Wiedemann, 1820, Nova Dipt. Gen.: 16. Type-species, *Graptomyza longirostris* Wiedemann, by present designation.

Baryterocera Walker, 1857, J. Linn. Soc. Lond. 1: 123. Type-species, *inclusa* Walker, by monotypy.

Ptilostylomyia Bigot, 1882, Annls Soc. ent. Fr. (6) 2 (Bull.): cxiv. Type-species, *Grap-*

myza brevirostris Wiedemann, by present designation.

References: Curran 1942; de Meijere 1908; Kertész 1914 (keys).

angustimarginata Brunetti. — India [Assam].

1923, Fauna Br. India, Dipt. 3: 141 (♀). Type-loc: Tura, Garo Hills, Assam, India.

arisana Shiraki. — Formosa.

1930, Mem. Fac. Agric. Taihoku imp. Univ. 1: 236 (♂). Type-loc: Arisan, Formosa.

brevirostris Wiedemann. — Borneo, Burma, Ceylon, India [Madras], Java, Malaya, Nepal, Philippines [Mindanao].

1820, Nova Dipt. Gen.: 17 (♀). Type-loc: Batavia (Djakarta), Java.

coomani Séguy. — Ceylon, N Viet Nam.

1948, Notes Ent. chin. 12: 159 (♂ ♀). Type-loc: Hoa-Binh, Tonkin, N Viet Nam.

cornuta de Meijere. — Java.

1914, Tijdschr. Ent. 57: 149 (♂). Type-loc: Gunung Gede, Java.

cynocephala Kertész. — Formosa.

1914, Annls hist.-nat. Mus. natn. hung. 12: 81 (♂). Type-loc: Chip-Chip, Formosa.

dentata Kertész. — Formosa.

1914, Annls hist.-nat. Mus. natn. hung. 12: 83 (♀). Type-locs: Chip-Chip and Toyenmongai, Formosa.

dolichocera Kertész. — Formosa.

1914, Annls hist.-nat. Mus. natn. hung. 12: 79 (♀). Type-locs: Kosempo, Chip-Chip, and Koshun, Formosa.

elegans Hull. — Java.

1950, Ann. Mag. nat. Hist. (12) 3: 619 (♀). Type-loc: Soekaboemi, Java.

fascipennis Sack. — Formosa.

1922, Arch. Naturgesch. 87 (11): 262 (♀). Type-loc: Shishastammes, Formosa.

flavipes de Meijere. — Borneo, Java, Philippines [Mindanao].

1911, Tijdschr. Ent. 54: 344 (♀). Type-loc: Gunung Ungaran, Java.

flavitincta Hull. — Java.

1950, Ann. Mag. nat. Hist. (12) 3: 622 (♀). Type-loc: Soekaboemi, Java.

flavonotata Brunetti. — India.

1917, Rec. Indian Mus. 13: 86 (♂). Type-loc: between Kufri and Phagu, Simla District, India.

flavorhyncha Hull. — Philippines [Mindanao].

1941, Psyche, Camb. 48: 158 (♂). Type-loc: Mainit River (6000 ft), Mt Apo, Mindanao, Philippines.

formosana Shiraki. — Formosa.

1930, Mem. Fac. Agric. Taihoku imp. Univ. 1: 234 (♂). Type-locs: Shu-Shu, Taihorin, Shinchiku, Arisan, and Kosempo, Formosa.

gibbula (Walker). — Celebes, Formosa.

1860, J. Linn. Soc. Lond. 4: 120 (♀; *Baryterocera*). Type-loc: Makasar, Celebes.

globigaster Hull. — Philippines [Biliran].

1943, J. Wash. Acad. Sci. 33: 42 (♀). Type-loc: Biliran Island, Philippines.

- hardyi* Greene. — India [Assam].
1949, Proc. ent. Soc. Wash. 51: 80 (♂). Type-loc: Misamari, Assam, India.
- inclusa* (Walker). — Borneo.
1857, J. Linn. Soc. Lond. 1: 123 (♀; *Baryterocera*). Type-loc: Sarawak, Borneo.
- interrupta* Wiedemann. — Java.
1830, Aussereurop. zweifl. Insekts. 2: 209 (♀). Type-loc: Java.
- jacobsoni* de Meijere. — Java, Sumatra.
1911, Tijdschr. Ent. 54: 343 (sex?). Type-locs: Telaga Mendjer and Gunung Ungaran, Java.
- javensis* Greene. — Java.
1949, Proc. ent. Soc. Wash. 51: 79 (♀). Type-loc: Mt Salak, Java.
- lineata* Osten Sacken. — Malaya; Moluccas.
1881, Annali Mus. civ. Stor. nat. Giacomo Doria 16: 439 (♀). Type-loc: Ternate, Moluccas.
- literata* Osten Sacken. — Philippines [Luzon].
1882, Berl. ent. Z. 26: 119 (? ♀). Type-loc: Philippines.
- longirostris* Wiedemann. — Burma, India [Assam], Java, Malaya, Sumatra.
1820, Nova Dipt. Gen.: 16 (? ♂). Type-loc: Java.
- ssp. *duodecimnotata* Brunetti. — India [Assam].
1908, Rec. Indian Mus. 2: 63 (sex?; as var. 12-notata). Type-loc: Sadiya, Assam, India.
- melliponaiformis* Doleschall. — Celebes; Moluccas (Amboina).
1858, Natuurk. Tijdschr. Ned.-Indië 17: 95 (sex?). Type-locs: Amboina, Moluccas and Makasar, Celebes.
- microdon* Osten Sacken. — Philippines [Luzon].
1882, Berl. ent. Z. 26: 119 (♂ ♀). Type-loc: Philippines.
- nigripes* Brunetti. — India [Assam, Sikkim, W Bengal], Malaya.
1913, Rec. Indian Mus. 8: 167 (♀; as *ventralis* var.). Type-loc: Sadiya, Assam, India.
- nitobei* Shiraki. — Formosa.
1930, Mem. Fac. Agric. Taihoku imp. Univ. 1: 238 (♂ ♀). Type-loc: Arisan, Formosa.
- obtusa* Kertész. — Formosa.
1914, Annls hist.-nat. Mus. natn. hung. 12: 77 (♀). Type-loc: Toyenmongai, Formosa.
- palmeri* Greene. — Java.
1949, Proc. ent. Soc. Wash. 51: 79 (♂). Type-loc: Mt Gede, Java.
- phyllocera* Hull. — Philippines [Luzon].
1950, Ann. Mag. nat. Hist. (12) 617 (♀). Type-loc: Baguio, Luzon, Philippines.
- punctata* de Meijere. — Java; New Guinea.
1908, Tijdschr. Ent. 51: 280 (sex?). Type-loc: Erima, Astrolabe-Bai, New Guinea.
- rectifacies* de Meijere. — Java, Sumatra.
1915, Tijdschr. Ent. 58 (Suppl.): 80 (♂ ♀). Type-locs: Matur, Sumatra; and Gunung Ungaran, Java.

- seimunda* Curran. — Malaya.
1928, J. fed. Malay St. Mus. 14: 173 (♀). Type-loc: Jor Camp (2000 ft), Perak, Malaya.
- setigloba* Hull. — Philippines [Mindanao].
1941, Psyche, Camb. 48: 159 (♂). Type-loc: Mainit River (6000 ft), Mt Apo, Mindanao, Philippines.
- sexnotata* Brunetti. — India [Assam, Sikkim].
1908, Rec. Indian Mus. 2: 63 (♀). Type-loc: Sikkim, India.
- spectralis* Hull. — Java.
1950, Ann. Mag. nat. Hist. (12) 3: 615 (♀). Type-loc: Soekaboemi, Java.
- tinctovittata* Brunetti. — India [W Bengal].
1915, Rec. Indian Mus. 11: 227 (? ♀). Type-loc: Sukna, Darjeeling Hills, India.
- ventralis* Wiedemann. — India [Assam], Java, Malaya.
1830, Aussereurop. zweifl. Insekts. 2: 207 (♀). Type-loc: Java.

Genus ORNIDIA Lepeletier and Serville

- Volucella*, subg. *Ornidia* Lepeletier and Serville, in Latreille, et al., 1828, Encycl. méth. (Ins.) 10: 786. Type-species, *Syphus obesus* Fabricius, by original designation.
- obesa* (Fabricius). — Java; throughout world tropics, Neotropical.
1775, Syst. Ent.: 763 (sex? *Syphus*). Type-loc: America.

Genus VOLUCELLA Geoffroy

- Volucella* Geoffroy, 1762, Hist. Insect. Paris 2: 540. Type-species, *Musca pellucens* Linnaeus, by designation of Curtis (1833: pl. 452). Validated by I.C.Z.N. 1957, Opinion 441: 85.
Reference: Coe 1964 (key, *bombylans* group).
- basalis* Brunetti. — India [Sikkim].
1907, Rec. Indian Mus. 1: pl. xi, figs. 11 & 12; 1908, Rec. Indian Mus. 2: 61 (♀). Type-loc: Mungphu, Sikkim, India.
- bombylans* (Linnaeus)
var. *malayana* Curran. — Malaya.
1928, J. fed. Malay St. Mus. 14: 321 (♂). Type-loc: Malay Peninsula.
- dimidiata* Sack. — Formosa.
1922, Arch. Naturgesch. (A) 87(11): 261 (♂). Type-loc: Taihorinsho, Formosa.
- discolor* Brunetti. — India [Assam]; Japan.
1908, Rec. Indian Mus. 2: 62 (♂). Type-loc: Sibsagar, Assam, India.
- flavolineata* Hull. — Philippines [Mindanao].
1943, Ann. Mag. nat. Hist. (11) 10: 29 (♀). Type-loc: Mindanao, Philippines.
- flavoscutella* Sack. — India [Uttar Pradesh].
1928, Entomologist's mon. Mag. 64: 107 (♀). Type-loc: Sangcha, N Kumaon, India.
- lividiventris* Brunetti. — India [Sikkim]; Tibet.
1908, Rec. Indian Mus. 2: 62 (♂ ♀). Type-loc: Sikkim, India.

- nigricans* Coquillett. — Formosa; Japan, Korea.
1898, Proc. U.S. natn. Mus. 21: 324 (♂ ♀). Type-loc: Japan.
- nigropictoides* Curran. — China [Fukien].
1925, Am. Mus. Novit. 200: 2 (♂ ♀). Type-loc: Yen-ping (Nanping), China.
- nitidithorax* Hull. — Philippines [Mindanao].
1941, Psyche, Camb. 48: 155 (♂). Type-loc: Mainit River (6500 ft), Mt Apo, Mindanao, Philippines.
- niveipes* Hull. — Java.
1943, Ann. Mag. nat. Hist. (11) 10: 27 (♂). Type-loc: Java.
- opalina* Wiedemann. — Not Oriental (see Therese, 1902, Berl. Ent. Ztg 47: 245); Neotropical.
- peleterii* Macquart. — India [Assam], Java.
1834, Hist. nat. Ins. Dipt. 1: 495 (♂). Type-loc: Java.
- signata* Brunetti, 1923, Fauna Br. India, Dipt. 3: 146 (♂). Type-loc: above Tura, Garo Hills, Assam, India.
- pellucens* (Linneaus). — India, Malaya; Europe, Japan.
1758, Syst. Nat. Ed. 10: 595 (sex? *Musca*). Type-loc: Europe.
- rotundata* Edwards. — Malaya, Sumatra.
1919, J. fed. Malay St. Mus. 8: 38 (♂). Type-loc: Sandaran Agong, Sumatra.
- ruficauda* Brunetti. — India [Sikkim].
1907, Rec. Indian Mus. 1: pl. xi, fig. 13; 1908, Rec. Indian Mus. 2: 61 (♂ ♀). Type-loc: Sikkim, India.
- surda* Curran. — Borneo, Thailand.
1928, J. fed. Malay St. Mus. 14: 163 (♀). Type-loc: Nakhon Si Thammarat, Khao Luang (2000 ft), Thailand.
- taiwana* Shiraki. — Formosa.
1930, Mem. Fac. Agric. Taihoku imp. Univ. 1: 215 (♂). Type-loc: Tenchosan-Karenko, Formosa.
- terauchii* Matsumura. — Formosa.
1916, Thousand Insects Japan 2: 237 (♀). Type-loc: Gyoichi nr Horisha, Formosa.
- trifasciata* Wiedemann. — Celebes, China, India to Formosa, Java, Philippines.
1830, Aussereurop. zweifl. Insekt. 2: 196 (♀). Type-loc: Java.
- linearis* Walker, 1852, Insecta Saundersiana 1: 251 (♀). Type-loc: unknown.
- decorata* Walker, 1860, J. Linn. Soc. Lond. 4: 120 (♂). Type-loc: Makasar, Celebes.
- nubeculosa* Bigot, 1875, Annls Soc. ent. Fr. 5(5): 474 (♀). Type-loc: China.
- var. *europila* Curran. — Malaya.
1928, J. fed. Malay St. Mus. 14: 163 (♀). Type-loc: Pahang, Cameron Highlands, No. 4 Camp (4800 ft), Malaya.
- ursina* de Meijere. — India [W Bengal].
1904, Bijdr. Dierk. 18: 101 (♀). Type-loc: Darjeeling, India.
- varipila* Coe. — India, Nepal.
1964, Bull. Br. Mus. nat. Hist. (Ent.) 15: 268 (♂ ♀). Type-loc: Khumbu, Khumdzung, E Nepal.
- vespimima* Shiraki. — Formosa.
1930, Mem. Fac. Agric. Taihoku imp. Univ. 1: 228 (♂ ♀). Type-loc: Arisan, Kobayashi, Karenko, Formosa.

- violacea* (Lepeletier and Serville). — China.
1828, Encycl. Méth. (Ins.) 10: 787 (♂; *Tenocera*). Type-loc: China.
- xanthopygata* Curran. — Java.
1928, J. fed. Malay St. Mus. 14: 168 (♂). Type-loc: Ongop Ongop, Java.

Tribe CHRYSOGASTERINI

Genus ORTHONEVRA Macquart

- Orthonevra* Macquart, 1829, Ins. Dipt. Nord Fr. 4: 40 (188). Type-species, *Chrysogaster elegans* Meigen, by monotypy.
Orthoneura, emendation.
- formosana* Shiraki. — Formosa..
1930, Mem. Fac. Agric. Taihoku imp. Univ. 1: 256 (♀). Type-loc: Toseikaku, Formosa.
- indica* Brunetti. — India [Punjab].
1915, Rec. Indian Mus. 11: 203 (♂). Type-loc: Kangra Valley, Punjab, India.

Genus MYOLEPTA Newman

- Myolepta* Newman, 1838, Ent. Mag. (Lond.) 5: 373. Type-species, *Musca luteola* Gmelin, by monotypy.
Myolepta, emendation.
- Reference: Thompson 1971 (rev.).
- himalayana* Brunetti. — India.
1915, Rec. Indian Mus. 11: 233 (♂ ♀). Type-loc: Matiana, Simla District, India.
- orientalis* Thompson. — Thailand.
1971, Proc. ent. Soc. Wash. 73: 346 (♀). Type-loc: Chiangdao (450 m), NW Chiangmai Province, Thailand.
- petiolata* Thompson. — Thailand.
1971, Proc. ent. Soc. Wash. 73: 344 (♀). Type-loc: Chiangdao (450 m), NW Chiangmai Province, Thailand.

Genus SPHEGINA Meigen

- Sphegina* Meigen, 1822, Syst. Beschr. europ. zweifl. Insekt. 3: 193. Type-species, *Milesia clunipes* Fallén, by designation of Westwood (1840: 136).

Subgenus SPHEGINA Meigen

- ?*asciiformis* Brunetti. — India [W Bengal].
1915, Rec. Indian Mus. 11: 223 (♀). Type-loc: Darjeeling, India.
- tricoloripes* Brunetti. — India [Himachal Pradesh, Uttar Pradesh].
1915, Rec. Indian Mus. 11: 225 (♀). Type-loc: Bhowali, Kumaon District, India.

Subgenus ASIOSPHEGINA Stackelberg

- Asiosphegina* Stackelberg, 1953, Trudy zool. Inst. Akad. Nauk. SSSR 13: 376.
Type-species, *Sphegina sibirica* Stackelberg, *in litt.*
- bispinosa* Brunetti. — India [Assam, Uttar Pradesh, W Bengal].
1915, Rec. Indian Mus. 11: 223 (♂ ♀). Type-locs: Margherita, Assam and Darjeeling, India.
- hansoni* Thompson. — Nepal.
1966, Bull. Brooklyn ent. Soc. 59/60: 42 (1964/1965) (♂). Type-loc: Parewair, Nepal.
- javana* de Meijere. — Java.
1914, Tijdschr. Ent. 57: 166 (♂). Type-loc: Gunung Gede, Java.
- ?*tenuis* Brunetti, 1915, Rec. Indian Mus. 11: 224 (♂, not ♀). Type-loc: Darjeeling, India.
- tristriata* Brunetti. — India [Assam].
1913, Rec. Indian Mus. 8: 165 (♀). Type-loc: Rotung, NE Frontier, Assam, India.

Unplaced species of *Sphegina* Meigen

- apicalis* Shiraki. — Formosa.
1930, Mem. Fac. Agric. Taihoku imp. Univ. 1: 423 (♀). Type-loc: Mu-sha, Formosa.
- nigerrima* Shiraki. — Formosa.
1930, Mem. Fac. Agric. Taihoku imp. Univ. 1: 425 (♂). Type-loc: Rantaian, Formosa.
- orientalis* Kertész. — Formosa, Philippines [Luzon].
1914, Annls hist.-nat. Mus. natn. hung. 12: 73 (♂ ♀). Type-locs: Chip-Chip, and Taihorin, Formosa.
- varidissima* Shiraki. — Formosa.
1930, Mem. Fac. Agric. Taihoku imp. Univ. 1: 421 (♀). Type-loc: Hori-sha, Formosa.

Tribe SPHEGINOBACCHINI

Genus SPHEGINOBACCHA de Meijere

- Spheginobaccha* de Meijere, 1908, Tijdschr. Ent. 51: 327. Type-species, *Sphegina macropoda* Bigot, by original designation.
- demeijerei* van Doesburg. — Java.
1968, Zoöl. Meded. Leiden 43: 161 (♂ ♀). Type-loc: Depok, Java.
- humeralis* (Sack). — Malaya.
1926, Philipp. J. Sci. 29: 571 (♂ ♀; Doros). Type-locs: Penang and Singapore, Malaya.
- liefitincki* van Doesburg. — Malaya.
1968, Zoöl. Meded. Leiden 43: 160 (♂ ♀). Type-loc: Sungai Raia, Ipoh, Perak, Malaya.

Syrphidae

macropoda (Bigot). — Anambas Islands, Borneo, Burma, Java, Laos, Malaya, Viet Nam.

1893, Annls Soc. ent. Fr. (6) 3: 331 (♀; *Sphegina*). Type-loc: Burma.
robusta Brunetti, 1907, Rec. Indian Mus. 1: pl. xi, figs. 3 & 4; 1908, Rec. Indian Mus. 2: 50 (♂ ♀; *Baccha*). Type-loc: Mergui, Burma.

melancholia Hull. — Viet Nam.

1937, J. Wash. Acad. Sci. 27: 174 (♀). Type-loc: Prang Bon, 30 mi NW of Saigon, Cochinchina.

Tribe SERICOMYINI

Genus PARARCTOPHILA Hervé-Bazin

Pararctophila Hervé-Bazin, 1914, Insecta 41: 152. Type-species, *oberthuri* Hervé-Bazin, by monotypy.

oberthuri Hervé-Bazin. — India [Uttar Pradesh], north to Siberia (Transbaikal) and east to China [Fukien, Hopeh, Kiangsu].

1914, Insecta 41: 153 (♂). Type-loc: "Padong, British Bootan, Inde."
simplicipes Brunetti, 1915, Rec. Indian Mus. 11: 247 (♀; *Arctophila*). Type-loc: Garhwal District, Kumaon, India.

Genus PSEUDOVOLOCCELLA Shiraki

Pseudovolucella Shiraki, 1930, Mem. Fac. Agric. Taihoku imp. Univ. 1: 39. Type-species, *mimica* Shiraki, by original designation.

Reference: Coe 1964 (key.).

apiformis (de Meijere). — Sumatra.

1919, Bijdr. Dierk. 21: 27 (♂; *Arctophila*). Type-loc: Piek von Korintji, Sumatra.

crassipes Edwards, 1919, J. fed. Malay St. Mus. 8: 44 (♂; *Sericomyia*). Type-loc: Korinchi Peak, Sumatra.

apimima Hull. — Java.

1941, Psyche, Camb. 48: 154 (♂). Type-loc: Tjibodas, Mt Gede, Java.

fasciata Curran. — Borneo.

1931, J. fed. Malay St. Mus. 16: 369 (♂ ♀). Type-loc: Mt Kinabalu, Lumu Lumu, N Borneo.

hingstoni Coe. — India [Sikkim], Nepal.

1964, Bull. Br. Mus. nat. Hist. (Ent.) 15: 270 (♂ ♀). Type-loc: Phadam Chen, Sikkim, India.

malayana (Curran). — Malaya.

1928, J. fed. Malay St. Mus. 14: 286 (♂; *Pararctophila*). Type-locs: Gunong Tahan (6500 ft) and Tanah Rata, Cameron Highlands, Pahang, Malaya.

mimica Shiraki. — Formosa.

1930, Mem. Fac. Agric. Taihoku imp. Univ. 1: 40 (♂ ♀). Type-locs: Asahi, and Tamaru, Formosa.

ochracea Hull. — Burma.

1944, Ann. Mag. nat. Hist. (11) 11: 32 (♂). Type-loc: Burma.

Genus SERICOMYIA Meigen

- Cinxia* Meigen, 1800, Nouv. Class. Mouches: 35. Type-species, *Musca lappona* Linnaeus, by designation of Coquillett (1910: 524). Suppressed by I.C.Z.N., 1963, Opinion 678: 339.
- Sericomyia* Meigen, 1803, Magazin Insektenkde 2: 274. Type-species, *Musca lappona* Linnaeus, by designation of Latreille (1810: 443).
- completa* Curran. — Formosa.
1929, Ann. Ent. Soc. Am. 22: 502 (♂). Type-loc: Kiluno, Formosa.
- cristaloides* Brunetti. — India [Assam].
1913, Rec. Indian Mus. 8: 167 (♀). Type-loc: between Renging and Rotun, NE Frontier, Assam, India.
- himalayensis* Brunetti. — India.
1907, Rec. Indian Mus. 1: pl. xiii, figs. 6–8; 1908, Rec. Indian Mus. 2: 89 (♂). Type-loc: Sikkim, India.

Tribe EUMERINI

Genus AZPEYTIA Walker

- Azpeytia* Walker, 1865, J. Linn. Soc. Lond. 8: 113. Type-species, *scutellaris* Walker, by monotypy.
- albomaculata* Curran. — Malaya.
1931, J. fed. Malay St. Mus. 16: 329 (♂). Type-loc: Bukit Kutu (3500 ft), Selangor, Malaya.
- bifascia* Brunetti. — India [Assam], Laos.
1907, Rec. Indian Mus. 1: pl. xiii, figs. 1–5; 1908, Rec. Indian Mus. 2: 84 (♂). Type-loc: Margherita, Assam, India.
- flavoscutellata* Kertész. — Formosa.
1913, Annls hist.-nat. Mus. natn. hung. 11: 284 (♂ ♀). Type-locs: Koso, and Mt Hoozan, Formosa.
- maculata* Shiraki. — Burma, Formosa.
1930, Mem. Fac. Agric. Taihoku imp. Univ. 1: 209 (♂ ♀). Type-locs: Karenko District, Formosa; and Tenasserim (Burma).
- scutellaris* Walker. — Thailand; New Guinea.
1865, J. Linn. Soc. Lond. 8: 113 (♂ ♀). Type-loc: New Guinea.
- trepidus* Curran, 1928, J. fed. Malay St. Mus. 14: 195 (♂; *Eumerus*). Type-loc: Khao Luang (2000 ft), Nakhon Si Thammarat, Thailand.

Genus EUMERUS Meigen

- Eumerus* Meigen, 1822, Syst. Beschr. europ. zweifl. Insekts. 3: 202. Type-species, *Syrphus tricolor* Fabricius, by designation of Curtis (1839: pl. 749).
- Citibaena* Walker, 1857, J. Linn. Soc. Lond. 1: 124. Type-species, *aurata* Walker, by monotypy.
- aeneithorax* Brunetti. — India.
1915, Rec. Indian Mus. 11: 244 (♂). Type-loc: Simla, India.

- albifrons* Walker. — India [Madras], Java.
1852, Insecta Saundersiana 1: 224 (♂). Type-loc: East Indies.
- halictiformis* Brunetti, 1915, Rec. Indian Mus. 11: 241 (♂ ♀). Type-loc: Orissa coast, Puri, India.
- argentipes* Walker. — Celebes, Ceylon, Malaya, Philippines [Mindanao]; Moluccas (Amboina, Batjan), New Guinea.
1861, J. Linn. Soc. Lond. 5: 284 (♀). Type-loc: Batjan, Moluccas.
- argyropus* Doleschall, 1857, Natuurk. Tijdschr. Ned.-Indië 14: 410 (sex?; preocc. Loew 1848). Type-loc: Amboina, Moluccas.
- doleschalli* Shiraki, 1930, Mem. Fac. Agric. Taihoku imp. Univ. 1: 97 (n. name for *argyropus* Doleschall).
- auratus* (Walker). — Borneo.
1857, J. Linn. Soc. Lond. 1: 124 (♂; *Citibaena*). Type-loc: Sarawak, Borneo.
- aurifrons* (Wiedemann). — Ceylon, Formosa, India [Bihar, Kashmir, Maharashtra], Java, Malaya, Philippines [Mindanao], Sumbawa; Australia.
1824, Analecta Ent.: 32 (♂; *Pipiza*). Type-loc: "Ostindien."
- var. *similis* Keiser. — Ceylon.
1958, Revue suisse Zool. 65: 216 (♀). Type-loc: Kandy, Deiyannewela, Ceylon.
- coeruleifrons* Keiser. — Ceylon.
1958, Revue suisse Zool. 65: 218 (♂). Type-loc: Peradeniya, Expt. Station, Ceylon.
- deceptor* Curran. — Borneo, Malaya.
1928, J. fed. Malay St. Mus. 14: 190 (♂ ♀). Type-locs: Selangor-Pahang border, Ginting Sempak and Perak, Batang Padang, Jor Camp (1800 ft), Malaya.
- ergator* Hull. — Philippines [Mindanao].
1941, Psyche, Camb. 48: 161 (♂). Type-loc: Mainit River (6500 ft), Mt Apo, Mindanao, Philippines.
- figurans* Walker. — Bali, Borneo, Celebes, Ceylon, Formosa, Java, Malaya, Philippines, Sumatra, Sumbawa; Hawaii.
1860, J. Linn. Soc. 4: 121 (♀). Type-loc: Makasar, Celebes.
- marginatus* Grimshaw, 1902, Fauna hawaiiensis 3: 82 (♂). Type-loc: Honolulu, Hawaii.
- flavicinctus* de Meijere. — Bali, Formosa, Java. Philippines [Luzon, Mindanao], Sumatra.
1908, Tijdschr. Ent. 51: 215 (♂ ♀). Type-locs: Semarang, Buitenzorg (Bogor), Java, and Medan, Fort de Kock, Sumatra.
- flavipes* Brunetti. — Borneo.
1915, Rec. Indian Mus. 11: 241 (♀). Type-loc: Kuching, Sarawak, Borneo.
- floresanus* Keiser. — W Flores.
1952, Verh. naturf. Ges. Basel 63: 166 (♂). Type-loc: Rana Mese, W Flores.
- halictoides* Brunetti. — India [W Bengal].
1915, Rec. Indian Mus. 11: 242 (♂ ♀). Type-locs: Darjeeling and Simla, India.

- insistens* Curran. — Malaya.
 1928, J. fed. Malay St. Mus. 14: 190 (♀). Type-loc: Lubok Kedondong, NW of Mt Ophir, Lalang and Beluka (200 ft), Malacca-Johore border, Malaya.
- macrocerus* Wiedemann. — China.
 1830, Aussereurop. zweifl. Insekts. 2: 113 (♂). Type-loc: China.
- nepalensis* Brunetti. — Borneo, Nepal.
 1908, Rec. Indian Mus. 2: 76 (♀). Type-loc: Chonebal, Nepal.
- nicobarensis* Schiner. — Borneo, Ceylon, Formosa, India [Assam, Kashmir, W Bengal], Malaya, Nicobar Islands.
 1868, in Reise der österreichischen Fregatte Novara, Dipt.: 368 (♀). Type-loc: Nicobar Islands.
- niveipes* de Meijere. — Java, Lesser Sunda Islands, Philippines [Luzon].
 1908, Tijdschr. Ent. 51: 220 (♂). Type-loc: Batavia (Djakarta), Java.
- okinawaensis* Shiraki. — Formosa, Ryukyu Islands (Okinawa); Japan.
 1930, Mem. Fac. Agric. Taihoku imp. Univ. 1: 98 (♂ ♀). Type-locs: Naha, Yayeyama, Haibara, Okinawa; Oshima, Kagoshima, Japan; and Kotosho, Formosa.
- parallelus* de Meijere. — Java.
 1908, Tijdschr. Ent. 51: 217 (♀). Type-loc: Batavia (Djakarta), Java.
- perpensus* Brunetti. — India.
 1917, Rec. Indian Mus. 13: 88 (♀). Type-loc: Phagu, Simla District, India.
- perplexus*, Brunetti. — India.
 1917, Rec. Indian Mus. 13: 89 (♀). Type-loc: Simla, India.
- pulcherrimus* Brunetti. — India [W Bengal].
 1915, Rec. Indian Mus. 11: 243 (♀). Type-loc: Kuresong, India.
- pulverulentus* Brunetti. — India [Kashmir, Rajasthan].
 1923, Fauna Br. India, Dipt. 3: 258 (♂ ♀). Type-loc: Pusa, India.
- quadratimaculatus* Keiser. — Sumbawa.
 1952, Verh. Naturf. Ges. Basel 63: 167 (♂). Type-loc: Baing, Sumbawa.
- rufitibiis* Sack. — Formosa.
 1922, Arch. Naturgesch. 87(11): 271 (♂ ♀; as *piveipes* var.). Type-loc: Kankau, Formosa.
- rufoscutellatus* Brunetti. — India [W Bengal], Thailand.
 1913, Rec. Indian Mus. 9: 269 (♂). Type-loc: Darjeeling, [W Bengal] India.
- sexvittatus* Brunetti. — India [Uttar Pradesh].
 1915, Rec. Indian Mus. 11: 245 (♀). Type-loc: Bhowali, Kumaon District, India.
- singhalensis* Keiser. — Ceylon.
 1958, Revue suiss Zool. 65: 219 (♂ ♀). Type-loc: Kalpitiya, Ceylon.
- sita* Keiser. — Ceylon.
 1958, Revue suisse Zool. 65: 217 (♀). Type-loc: Kandy, Hantana, Ceylon.
- splendens* Wiedemann. — Borneo, ?Java.
 1830, Aussereurop. zweifl. Insekts. 2: 114 (♀). Type-loc: East Indies.

sutteri Keiser. — Sumbawa.

1952, Verh. Naturf. Ges. Basel 63: 168 (♂ ♀). Type-loc: Waimangura, Matakori, W Sumbawa.

Genus MERODON Meigen

- Lampetia* Meigen, 1800, Nouv. Class. Mouches: 34. Type-species, *Syrphus clavipes* Fabricius, by designation of Coquillett (1910: 557). Suppressed by I.C.Z.N. 1963, Opinion 678: 339.
- Merodon* Meigen, 1803, Magazin Insektskde 2: 274. Type-species, *Syrphus clavipes* Fabricius, by designation of Westwood (1840: 137) (as *curvipes*, error by Meigen).
- albifasciatus* Macquart. — India or Java.
 1842, Dipt. exot. 2(2): 71 (131) (♀). Type-loc: "Indes Orientales."
- pruni* (Rossi). — W Pakistan; Palaearctic.
 1790, Fauna Etrusca 2: 293 (sex?). Type-loc: Italy.
- pallidus* Macquart, 1842, Dipt. exot. 2(2): 70 (130) (♀). Type-loc: Bagdad, Iraq.
- tuberculatus* (Brunetti). — India [W Bengal].
 1907, Rec. Indian Mus. 1: pl. xii, figs. 16 & 17; 1908, Rec. Indian Mus. 2: 67 (♂ ♀; *Helophilus*). Type-loc: Calcutta, India.
- varicolor* Walker. — Borneo, India [Assam].
 1857, J. Linn. Soc. Lond. 1: 122 (♀). Type-loc: Sarawak, Borneo.

Genus PSILOTA Meigen

- Psiolota* Meigen, 1822, Syst. Beschr. europ. zweifl. Insekts. 3: 256. Type-species, *anthracina* Meigen, by monotypy.
- bifenestrata* de Meijere. — Java.
 1933, Tijdschr. Ent. 76: 104 (♂ ♀). Type-locs: Parengan, Bodjonegoro, Semarang, and Krawang, Java.
- fasciata* Curran. — Philippines [Luzon].
 1929, Ann. Ent. Soc. Am. 22: 494 (♂). Type-loc: Manila, Luzon, Philippines.
- tectonae* de Meijere. — Java.
 1933, Tijdschr. Ent. 76: 103 (♂). Type-locs: Semarang, and Parengan, Java.

Tribe CERIOLIDINI

Genus CERIANA Rafinesque

- Ceria* Fabricius, 1794, Ent. Syst. 4: 277 (preocc. Scopoli 1763). Type-species, *clavicornis* Fabricius, by subsequent designation of Latreille (1810: 443), = *conopsoides* (Linnaeus).
- Cina* Fabricius, 1798, Suppl. ent. Syst.: 557. Misspelling.
- Ceriana* Rafinesque, 1815, Analyse: 131 (n. name for *Ceria* Fabricius). Type-species, *Ceria clavicornis* Fabricius, automatic, = *conopsoides* (Linnaeus).
- Tenthredomyia* Shannon, 1925, Insecutor Inscit. menstr. 13: 50. Type-species, *Ceria abbreviata* Loew, by original designation.

References: de Meijere 1908; Kertész 1913; Shannon 1927 (keys); Vockeroth 1971 (nomencl.); Bhatia 1931 (biol., imm., stages).
anaglypha (Séguy). — China.

- 1948, Notes ent. chin. 12: 163 (♂ ♀; *Ceriodes*). Type-loc: Kou-ling, China. N. comb.
brevis (Brunetti). — India [Bihar, Kashmir].
 1923, Fauna Br. India, Dipt. 3: 328 (♀; *Ceria*). Type-loc: Pusa, India. N. comb.
compacta (Brunetti). — India [Uttar Pradesh].
 1907, Rec. Indian Mus. 1: pl. xiii, fig. 13; 1908, Rec. Indian Mus. 2: 95 (♀; *Ceria*). Type-loc: Mussoorie, W Himalayas, India.
dimidiatispennis (Brunetti). — India [Kashmir], W Pakistan.
 1923, Fauna Br. India, Dipt. 3: 329 (♂ ♀; *Ceria*). Type-locs: Hangu, NW Frontier Province and Abbottabad, NW India (W Pakistan). N. comb.
formosensis (Shiraki). — Formosa.
 1930, Mem. Fac. Agric. Taihoku imp. Univ. 1: 3 (♂; *Cerioides*). Type-loc: Musha and Shinnensho, Formosa.
ornatifrons (Brunetti). — Ceylon, India [W Bengal], Nepal.
 1915, Rec. Indian Mus. 11: 252 (♀; *Ceria*). Type-loc: Kumdhik, Himalayas, Nepal.

Genus MONOCEROMYIA Shannon

Cerioides, subg. Monoceromyia Shannon, 1922, Bull. Brooklyn ent. Soc. 17: 32. Type-species, *Ceria tricolor* Loew, by monotypy.

References: see under *Ceriana*.

- anchorata* (Bigot). — Borneo.
 1883, Annls Soc. ent. Fr. (6) 3: 318 (♂ ♀; *Sphyximorpha*). Type-loc: Sarawak, Borneo.
annulata (Kertész). — Formosa; Japan.
 1913, Annls hist.-nat. Mus. natn. hung. 11: 407 (♂ ♀; *Cerioides*). Type-locs: Fuhosho, Suihenkyaku, Kanshirei, and Toyenmongai, Formosa.
horishana Matsumura, 1916, Thousand Insects Japan 2: 186 (♂; *Sphyximorpha*). Type-loc: Horisha, Formosa.
crux (Brunetti). — India [Uttar Pradesh].
 1915, Rec. Indian Mus. 11: 253 (♀; *Ceria*). Type-loc: Kousanie, Kumaon, W Himalayas, India.
eumenioides (Saunders). — India [Madras].
 1841, Trans. ent. Soc. Lond. 3: 60 (sex?; *Ceria*). Type-loc: northern India.
apicata Brunetti, 1908, Rec. Indian Mus. 2: 95 (♂; *Ceria*). Type-loc: Ganjam, Madras, India.
fenestrata (Brunetti). — India [Sikkim].
 1923, Fauna Br. India, Dipt. 3: 333 (♀; *Ceria*). Type-loc: Ronjit Valley, Sikkim, India.
flavipennis (de Meijere). — Celebes, India.
 1908, Tijdschr. Ent. 51: 195 (♂; *Cerioides*). Type-loc: Minahassa, Celebes.
himalayensis (de Meijere). — India [Sikkim].
 1908, Tijdschr. Ent. 51: 198 (♀; *Cerioides*). Type-loc: Sikkim, India.
javana (Wiedemann). — Borneo, India [Assam, Mysore, W Bengal], Java, Malaya, Nepal, Sumatra.
 1824, Analecta Ent. 32 (♀; *Ceria*). Type-loc: Java.

- lateralis* (Walker). — Celebes.
 1860, J. Linn. Soc. Lond. 4: 118 (♂; *Ceria*). Type-loc: Makasar, Celebes.
multipunctata (Hull). — India.
 1941, Psyche, Camb. 48: 163 (♂; *Cerioides*). Type-loc: Koolos, India. N. comb.
obscura (Brunetti). — India [Sikkim], Nepal.
 1907, Rec. Indian Mus. 1: pl. xiii, fig. 12; 1908, Rec. Indian Mus. 2: 94 (♀; *Ceria*). Type-loc: Sikkim, India.
patricia (Brunetti). — Burma, India.
 1923, Fauna Br. India, Dipt. 3: 335 (♂ ♀; *Ceria*). Type-loc: Tavoy, Tenasserim, Lower Burma.
petersi (Speiser). — Philippines [Luzon].
 1924, Wien. ent. Ztg 41: 53 (♀; *Cerioides*). Type-loc: Philippines.
bakeri Shannon, 1925, Insecutor Inscit. menstr. 13: 55 (sex?). Type-loc: Santa Maria, Laguna, Luzon, Philippines.
polistoides (Brunetti). — India, Nepal.
 1923, Fauna Br. India, Dipt. 3: 335 (♀; *Ceria*). Type-loc: Simla, India.
salvazai Shannon. — Laos.
 1927, J. Wash. Acad. Sci. 17: 47 (♂). Type-loc: Ban Sen Savouane, Luang Prabang, Laos.
siamensis (Curran). — Thailand.
 1931, J. fed. Malay St. Mus. 16: 299 (♀; *Cerioides*). Type-loc: Nan, Thailand.
similis (Kertész). — Formosa.
 1913, Annls hist.-nat. Mus. natn. hung. 11: 405 (♂ ♀; *Cerioides*). Type-locs: Kosempo, Fuhosho, and Toyenmongai, Formosa.
stenogaster Séguy. — Malaya.
 1948, Notes ent. chin. 12: 162 (♀). Type-loc: Penang, Malacca, Malaya.
tredecimpunctata (Brunetti). — Burma, India [Assam], Indochina.
 1923, Fauna Br. India, Dipt. 3: 336 (♂ ♀; *Ceria*). Type-locs: Salween Valley, Burma; and Lower Ranges, Khasi Hills, Assam, India.
trinotata (de Meijere). — Burma, India [W Bengal], Indochina, Laos, Malaya.
 1904, Bijdr. Dierk. 18: 97 (♂; *Ceria*). Type-loc: Darjeeling, India.
wallacei Shannon. — Celebes.
 1927, J. Wash. Acad. Sci. 17: 47 (♀). Type-loc: Makasar, Celebes.
wiedemanni Shannon. — Laos.
 1927, J. Wash. Acad. Sci. 17: 47 (♂). Type-loc: Ban Nam Mo, Luang Prabang, Laos.

Genus PRIMOCERIOIDES Shannon

- Primoceroides* Shannon, 1927, J. Wash. Acad. Sci. 17: 41. Type-species, *Cerioides petri* Hervé-Bazin, by original designation.
yoshikawai (Sasakawa). — Laos.
 1960, Nature & Life S.E. Asia 1: 449 (♂; *Cerioides*). Type-loc: Thakhek, Laos.

Genus SPHIXIMORPHA Rondani

- Sphiximorpha* Rondani, 1850, Annls Soc. ent. Fr. (2) 8: 212. Type-species, *Ceria subsessilis* Illiger, by original designation.
- Cerioides* Rondani, 1850, Annls Soc. ent. Fr. (2) 8: 211. Name cited in synonymy of *Sphiximorpha* Rondani. Validated by subsequent usage by Becker, et al., 1907: 156.
- Polybiomyia* subg. *Ceriatrichia* Hull, 1949, Trans. zool. Soc. Lond. 26: 381. Type-species, *Cerioides bulbosa* de Meijere, by original designation.
- References: see under *Ceriana*.
- bulbosa* (de Meijere). — Sumatra.
1924, Tijdschr. Ent. 67 (Suppl.): 18 (♂; *Cerioides*). Type-loc: Aur Kumanis, Sumatra.
- decorata* (Brunetti). — Burma.
1923, Fauna Br. India, Dipt. 3: 326 (♂; *Ceria*). Type-loc: Rangoon, Burma.
- fruhstorferi* (de Meijere). — India [Sikkim].
1908, Tijdschr. Ent. 51: 196 (♀; *Cerioides*). Type-loc: Sikkim, India.
- fulvescens* (Brunetti). — India [Uttar Pradesh].
1915, Rec. Indian Mus. 11: 251 (♂; *Ceria*). Type-loc: Bhowali, Kumaon District, W Himalayas, India.
- meijerei* (Shannon). — Indochina.
1927, J. Wash. Acad. Sci. 17: 45 (♀; *Cerioides*). Type-loc: Tong Lap, Haut Mekong, Indochina.
- polista* (Séguy). — Laos.
1948, Notes ent. chin. 12: 161 (♂; *Ceriana*). Type-loc: Nam-lang, Laos.
- triangulifera* (Brunetti). — India [W Bengal].
1913, Rec. Indian Mus. 9: 273 (♂ ♀; *Ceria*). Type-loc: Darjeeling District, India.

Tribe ERISTALINI

Genus AXONA Walker

- Axona* Walker, 1864, J. Linn. Soc. Lond. 7: 211. Type-species, *volucelloides* Walker, by monotypy, = *chalcopyga* (Wiedemann).
- chalcopyga* (Wiedemann). — Borneo, Burma, Indochina, Laos, Malaya, Philippines [Luzon, Sibuan, Mindanao, Basilan], Thailand; Guadalcanal, Micronesia, Moluccas (Amboina, Mysol).
1830, Aussereurop. zweifl. Insekt. 2: 178 (♂; *Eristalis*). Type-loc: Manila, Philippines.
- maxima* Doleschall, 1857, Natuurk. Tijdschr. Ned.-Indië 14: 405 (sex?; *Eristalis*). Type-loc: Amboina, Moluccas.
- transpositus* Walker, 1860, Trans. ent. Soc. Lond. (n. Ser.) 5: 289 (♂ ♀; *Eristalis*). Type-loc: Burma.
- volucelloides* Walker, 1864, J. Linn. Soc. Lond. 7: 212 (♂). Type-loc: Mysol, Moluccas.

Genus CATACORES Hull

- Catacores* Hull, 1944, Ent. News 55: 205. Type-species, *Axona cyanea* Brunetti, by original designation.
- cyanea* (Brunetti). — India [W Bengal].
1913, Rec. Indian Mus. 9: 272 (♂; *Axona*). Type-loc: Singla, Darjeeling, India.

= *Pseudostethus* (?)
(Kerteszonyi)

Genus DIGULIA de Meijere

- Digulia* de Meijere, 1913, Nova Guinea 9: 357. Type-species, *kochi* de Meijere, by monotypy.
- kochi* de Meijere. — Ceylon, Java, Karimun Islands, New Guinea.
1913, Nova Guinea 9: 357. (♀). Type-loc: Digul River, New Guinea.

Genus DISSOPTERA Edwards

- Dissoptera* Edwards, 1915, Trans. zool. Soc. Lond. 20: 410. Type-species, *pollinosa* Edwards, by monotypy.
- heterothrix* (de Meijere). — Borneo; Australia, New Guinea.
1908, Tijdschr. Ent. 51: 273 (♂ ♀; *Eristalis*). Type-locs: Insel Cretin (Tami); Mahakkam, Borneo.
- ?*pollinosa* Edwards, 1915, Trans. zool. Soc. Lond. 20: 410 (♂ ♀). Type-loc: Iwaka River, New Guinea.

Genus ERISTALINUS Rondani

- Eristalinus* Rondani, 1845, Nuovi Annali Sci. nat. Bologna 2(2): 453. Type-species, *Musca sepulchralis* Linnaeus, by designation of Rondani (1857: 38).
- Lathyrophthalmus* Mik, 1897, Wien. ent. Ztg 16: 114. Type-species, *Conops aeneus* Scopoli, by original designation.
- References: Hervé-Bazin 1923; Curran 1947 (keys); Metcalf 1913 (biol., imm. stages).
- aeneus* (Scopoli). — northern Oriental region; Australia, Bermuda, Hawaii, North America, Palaearctic.
1763, Ent. Carniolica: 356 (sex?; *Conops*). Type-loc: Europe. N. comb.
- var. *nigrolineatus* (Hervé-Bezin). — W Pakistan.
1923c, Annls Sci. nat. (10) 6: 134 (♂ ♀; *Lathyrophthalmus*). Type-loc: Kurrachee (Karachi?). N. comb.
- var. *taphicus* (Wiedemann). — China, India [Maharashtra, Uttar Pradesh], W Pakistan; Cyprus, Egypt, France, Korea, Monaco.
1830, Aussereurop. zweifl. Insekt. 2: 191 (♀; *Eristalis*; as sp.). Type-loc: Egypt. N. comb.
- arvorum* (Fabricius). — entire SE Asia; Australia, Hawaii, Japan, Micronesia.
1787, Mantissa Insect. 2: 335 (♂ ♀; *Syrphus*). Type-loc: China. N. comb.
- quadrilineatus* Fabricius, 1787, Mantissa Insect. 2: 336 (sex?; *Syrphus*). Type-loc: Tranquebar, India.
- tranquebarica* Gmelin, 1790, Syst. Nat. 5: 2870 (sex?; *Musca*; unjustified n. name for *quadrilineatus* Fabricius).
- fulvipes* Macquart, 1846, Dipt. exot. Suppl. 1: 128 (256) (♂; *Eristalis*). Type-loc:

- Australia.
- antidotus* Walker, 1849, List Dipt. Colln Br. Mus. 3: 626 (♂; *Eristalis*). Type-loc: China.
- anicetus* Walker, 1849, List Dipt. Colln Br. Mus. 3: 624 (♂; *Eristalis*). Type-loc: unknown.
- fo* Bigot, 1880, Annls Soc. ent. Fr. (5) 10: 220 (♂; *Eristalomyia*). Type-loc: Amoy, China.
- cunotata* Bigot, 1891, Nouv. Archs Mus. Hist. nat. Paris (3) 2: 208 (♂; *Eristalomyia*). Type-loc: Laos.
- cinereus* (de Meijere). — Sumatra.
1924, Tijdschr. Ent. 67 (Suppl.): 20 (♀; *Eristalis*). Type-loc: Serapai, Korintji, Sumatra. N. comb.
- connectens* (Hervé-Bazin). — India, Java, Viet Nam.
1923, Annls Sci. nat. (10) 6: 148 (♂ ♀; *Lathyrophthalmus*). Type-loc: Saigon, Indochina. N. comb.
- haileyburyi* (Nayar). — India [Uttar Pradesh].
1968, Pan-Pacif. Ent. 44: 121 (♀; *Lathyrophthalmus*). Type-loc: St John's College, Agra, India. N. comb.
- invirgulatus* (Keiser). — Ceylon.
1958, Revue suisse Zool. 65: 231 (♂; *Lathyrophthalmus*). Type-loc: Kandy, Hantana, Ceylon. N. comb.
- kochi* (de Meijere). — Sumbawa; Australia, New Guinea.
1908, Tijdschr. Ent. 51: 255 (♂ ♀; *Eristalis*). Type-loc: Etna-Bai, Merake, S New Guinea. N. comb.
- laetus* (Wiedemann). — Ceylon, China, Formosa, India [Andhra Pradesh, Maharashtra], Java, Malaya, Nepal, Philippines [Luzon], Sumatra, Sumbawa.
1830, Aussereurop. zweifl. Insekts. 2: 192 (♂; *Eristalis*). Type-loc: China. N. comb.
- pallinevis* Macquart, 1842, Dipt. exot. 2(2): 46 (106) (♂; *Eristalis*). Type-loc: Bengal, India. N. synonym.
- obscuritarsis* de Meijere, 1908, Tijdschr. Ent. 51: 250 (♂ ♀; *Eristalis*). Type-locs: Semarang, Java, Singapore, Malaya; and Bombay, India. N. synonym.
- lalitai* (Nayar). — India [Uttar Pradesh].
1968, Pan-Pacif. Ent. 44: 119 (♀; *Lathyrophthalmus*). Type-loc: St. John's College, Agra, India. N. comb.
- lucilia* (de Meijere). — Ceylon, Java, Simeuluë Island, Sumatra.
1911, Tijdschr. Ent. 54: 341 (♂; *Eristalis*). Type-loc: Semarang, Java. N. comb.
- lugens* (Wiedemann). — China, Formosa.
1830, Aussereurop. zweifl. Insekts. 2: 193 (♂; *Eristalis*). Type-loc: China. N. comb.
- albitarsis* Sack, 1917, Stettin. ent. Ztg 88: 312 (♂ ♀; *Lathyrophthalmus*). Type-locs: Suisharyo, and Chosokai, Formosa. N. synonym.
- nigroscutatus* (de Meijere). — Borneo, Indochina, Java, Simeuluë Island, Sumatra, Sumbawa.
1911, Tijdschr. Ent. 54: 337 (♂; *Eristalis*). Type-locs: Tandjong Priok, and Muara Angkee nr Batavia (Djakarta), Java. N. comb.
- nitidus* (van der Wulp). — Borneo, Indochina, Java, Sumatra; Fiji, Samoa.
1884, Annls Soc. ent. Belg. 28 (Bull.): ccxci (♀; *Eristalis*). Type-loc: Java. N. comb.

- obliquus* (Wiedemann). — southern Oriental region; New Guinea.
1824, Analecta Ent.: 38 (♂ ♀; *Eristalis*). Type-loc: Bengal, India. N. comb.
- obtusus* (Hull). — Java.
1941, Psyche, Camb. 48: 153 (♂; *Lathyrophthalmus*). Type-loc: Buitenzorg (Bogor), Java. N. comb.
- quadristriatus* (Macquart). — Ceylon, Formosa, India, Nepal, Philippines [Negros].
1846, Dipt. exot. Suppl. 1: 127 (245) (♀; *Eristalis*). Type-loc: "Inde." N. comb.
- quinquelineatus* (Fabricius). — Not Oriental.
- quinquestriatus* (Fabricius). — throughout Oriental region; Japan.
1794, Ent. Syst. 4: 289 (sex?; *Syrphus*). Type-loc: "India orientali." N. comb.
- aesepus* Walker, 1849, List Dipt. Colln Br. Mus. 3: 625 (♂ ♀; *Eristalis*). Type-loc: China.
- picta* Bigot, 1880, Annls Soc. ent. Fr. (5) 10: 219 (♂; *Eristalomyia*). Type-loc: "Indostan."
- rufus* (van der Goot). — Indochina, Java.
1964, Beaufortia 10: 220 (*Eristalis*; n. name for *ferrugineus* de Meijere). *ferrugineus* de Meijere, 1911, Tijdschr. Ent. 54: 339 (♂ ♀; *Eristalis*); preocc. Fabricius 1805). Type-loc: Muara Antjol, nr Batavia (Djakarta) and Pasuruan, Java.
- selectans* (Curran). — Malaya.
1931, J. fed. Malay St. Mus. 16: 334 (♂; *Lathyrophthalmus*). Type-loc: Bukit Kutu (3500 ft), Selangor, Malay. N. comb.
- sepulchralis* (Linnaeus). — China [Chekiang, Kiangsu], India; Europe.
1758, Syst. Nat. Ed. 10: 596 (sex?; *Musca*). Type-loc: Europe.
- sinabangensis* (de Meijere). — Simeuluë Island.
1915, Tijdschr. Ent. 58 (Suppl.): 31 (♀; *Eristalis*). Type-loc: Sinabang, Simalur.
- suavissimus* (Walker). — Philippines [Luzon]; Moluccas (Aru), Australia, Guadalcanal, Micronesia, New Guinea.
1859, J. Linn. Soc. Lond. 3: 95 (♀; *Eristalis*). Type-loc: Aru Islands, Moluccas. N. comb.
- tarsalis* (Macquart). — China, Formosa, India, Nepal; Japan, Korea.
1855, Dipt. exot. Suppl. 5: 87 (107) (♂ ♀; *Eristalis*). Type-loc: "Chine boreale." N. comb.
- ocularius* Coquillett, 1898, Proc. U.S. natn. Mus. 21: 325 (♂ ♀; *Eristalis*). Type-loc: Japan. N. synonym.
- tristriatus* (de Meijere). — Java, Philippines [Negros].
1911, Tijdschr. Ent. 54: 342 (♂; *Eristalis*). Type-locs: Semarang and Batavia (Djakarta), Muara Angke, Java. N. comb.
- Subgenus ERISTALODES Mik
- Eristalodes* Mik, 1897, Wien. ent. Ztg 16: 114. Type-species, *Eristalis taeniops* Wiedemann, by original designation.
- paria* (Bigot). — Ceylon, Formosa, India [Himachal Pradesh, Madras, Mysore, Uttar Pradesh], Java, Laos, Malaya, Nepal; Japan, Moluccas (Ternate).
1880, Annls Soc. ent. Fr. (5) 10: 218 (♂ ♀; *Eristalomyia*). Type-loc: Cey-

ion. N. comb.

zebrina Bigot, 1880, Annls Soc. ent. Fr. (5) 10: 222 (♂; *Eristalomyia*). Type-loc: Ternate, Moluccas.

kobusi de Meijere, 1908, Tijdschr. Ent. 51: 252 (♂ ♀; *Eristalis*). Type-loc: Tosari, W Java.

arisanus Matsumura, 1916, Thousand Insects Japan 2: 264 (♂; *Eristalis*). Type-loc: Arisan, Formosa.

orientalis Brunetti, 1923, Fauna Br. India, Dipt. 3: 183 (♂ ♀; as *Eristalis quinquelineatus* var.). Type-loc: India.

taeniops (Wiedemann). — India, Nepal, W Pakistan; Africa, Canary Islands, Palearctic.

1818, Zool. Mag. (Wied.) 2: 42 (sex?; *Eristalis*). Type-loc: "Cape." N. comb.

Subgenus MERODONOIDES Curran

Merodonoides Curran, 1931, J. fed. Malay St. Mus. 16: 333. Type-species, *circularis* Curran, by original designation, = *multifarius* (Walker).

multifarius (Walker). — Ceylon, India [Assam, Madhya Pradesh, Madras], Java, Malaya, Nepal, Viet Nam.

1852, Insecta Saundersiana 1: 248 (♀; *Eristalis*). Type-loc: East Indies. N. comb.

circularis Curran, 1931, J. fed. Malay St. Mus. 16: 333 (♂; *Merodonoides*). Type-loc: Kedah Peak (3300 ft), Malaya.

minutus Hull, 1944, Ann. Mag. nat. Hist. (11)11: 43 (♂; *Merodonoides*). Type-loc: Jubbulpore, India.

czernyi Hull, 1944, J. Wash. Acad. Sci. 34: 400 (♂; *Merodonoides*). Type-loc: Montes Mauson, Tonkin, N Viet Nam. N. synonym.

kandyensis Keiser, 1958, Revue suisse Zool. 65: 234 (♂; *Eristalis*). Type-loc: Kandy, Asgiriya Ceylon. N. synonym.

Genus ERISTALIS Latreille

Tubifera Meigen, 1800, Nouv. Class. Mouches: 34. Type-species, *Musca tenax* Linnaeus, by designation of Coquillett (1910: 618). Suppressed by I.C.Z.N. 1963, Opinion 678: 339.

Elophilus Meigen, 1803, Magazin Insektenkde 2: 274. Type-species, *Musca tenax* Linnaeus (as *Eristalis tenax* Fabricius), by designation of Latreille (1810: 443). (see Stone, et al. 1965: 622).

Eristalis Latreille, 1804, Nouv. Dict. Hist. nat. 24 (Tab. méth. Ins.): 194. Type-species, *Musca tenax* Linnaeus, by designation of Curtis (1832: pl. 432).

Eristaloidea Rondani, 1845, Nuovi Annali Sci. nat. Bologna 2(2): 453. Type-species, *Musca tenax* Linnaeus, by designation of Coquillett (1910: 540).

Eristalomyia Rondani, 1857, Dipterol. ital. Prodr. 2: 38. Type-species, *Musca tenax* Linnaeus, by original designation.

Eristalomyia, error or emendation.
References: Trybom 1875; Gabler 1932; Hartley 1961, 1963 (biol. imm. stages).

Subgenus ERISTALIS Latreille

proserpina Wiedemann. — China.

1830, Aussereurop. zweifl. Insekts. 2: 157 (♂). Type-loc: China.

tenax (Linnaeus). — entire Oriental region; cosmopolitan.

1758, Syst. Nat. Ed. 10: 591 (sex?; *Musca*). Type-loc: Europe.

cognatus Wiedemann, 1824, Analecta Ent.: 37 (♂). Type-loc: Tranquebar, India. N. synonym.

sinensis Wiedemann, 1824, Analecta Ent.: 37 (♂). Type-loc: China. N. synonym.

ventralis Thomson, 1869, in K. svenska fregatten Eugenies resa, Zool., Dipt.: 489 (♂ ♀). Type-loc: China. N. synonym.

Subgenus EOSERISTALIS Kanervo

Eoseristalis Kanervo, 1938, Annls Univ. fenn. abo. (A) 6: 12. Type-species, *Eristalis cerealis* Fabricius, by original designation.

albibasis Bigot. — India [Himachal Pradesh]; China [Szechwan].

1880, Annls Soc. ent. Fr. (5) 10: 215 (♀). Type-loc: "Indostan."

arbustorum (Linnaeus). — India [Kashmir]; Holarctic.

1758, Syst. Nat. Ed. 10: 591 (sex?; *Musca*). Type-loc: Europe.

bicornuta de Meijere. — Java.

1924, Tijdschr. Ent. 67: 216 (♂ ♀). Type-loc: Pangerango, W Java.

bidentata Sack. — Philippines [Mindanao, Calicoan].

1926, Philipp. J. Sci. 29: 579 (♂). Type-locs: Kolambungan, Lanao, and Davao, Mindanao, Philippines.

brevifacies Coe. — India, Nepal.

1964, Bull. Br. Mus. nat. Hist. (Ent.) 15: 274 (♂ ♀). Type-loc: between Sangu and Tamrang, Taplejung District, E Nepal.

cerealis Fabricius. — throughout Oriental region; Japan, Korea, Sakhalin.

1805, Syst. Antiliat.: 232 (sex?). Type-loc: China.

solutus Walker, 1849, List. Dipt. Colln Br. Mus. 3: 619 (♂). Type-loc: Nepal.

incisuralis Loew, 1858, Wien. ent. Mschr. 2: 108 (♂ ♀). Type-loc: Japan.

barbata Bigot, 1880, Annls Soc. ent. Fr. (5) 10: 214 (♂). Type-loc: "Indostan."

cerealoides Curran. — Borneo.

1931, J. fed. Malay St. Mus. 16: 374 (♀). Type-loc: Kamborangah, N Borneo.

curvipes Schiner. — Ceylon.

1868, in Reise der österreichischen Fregatte Novara, Dipt. 2: 363 (♂ ♀). Type-loc: Ceylon.

himalayensis Brunetti. — Burma, China [Szechwan], India [Himachal Pradesh, Kashmir, Sikkim, Uttar Pradesh, W Bengal], Nepal.

1908, Rec. Indian Mus. 2: 70 (n. name for *ursinus* Bigot).

ursinus Bigot, 1880, Annls Soc. ent. Fr. (5) 10: 215 (♂; preocc. Jaennicke 1867). Type-loc: "Indostan."

horticola (De Geer). — India, ?Malaya; Europe, N Africa.

1776, Mém. pour serv. Hist. Ins. 6: 140 (sex?; *Musca*). Type-loc: Europe.

basifemorata Brunetti, 1923, Fauna Br. India, Dipt. 3: 175 (♀). Type-loc: India.

intricariooides Brunetti. — India [Sikkim], Nepal.

1923, Fauna Br. India, Dipt. 3: 171 (♀). Type-loc: Sikkim, India.

simplicipes Curran. — Malaya.

1928, J. fed. Malay St. Mus. 14: 300 (♂ ♀). Type-locs: Gunung Tahan (7186 ft), Gunung Gedong (6400 ft), Rhododendron Hill (5200 ft), Cameron Highlands, and Lubok Tamang (3500 ft), Pahang, Malaya.

suturalis Brunetti. — India [Assam], Laos, Viet Nam.

1923, Fauna Br. India, Dipt. 3: 193 (♂ ♀). Type-loc: Khasi Hills, Assam, India.

vitalisi Hervé-Bazin, 1923, Bull. Soc. ent. Fr. 1923: 26 (♂ ♀). Type-locs: Laos; and Tonkin, Viet Nam.

Unplaced species of *Eristalis* Latreille

aequipars Walker. — Moluccas (Mysol).

1864, J. Linn. Soc. Lond. 7: 210 (♂). Type-loc: Mysol, Moluccas.

angustimarginalis Brunetti. — Burma, India [Assam].

1923, Fauna Br. India, Dipt. 3: 176 (♂ ♀). Type-locs: Takepum Mt, Chinese Frontier, NE Burma; and Shillong, India.

cingulata Sack. — Formosa, Philippines [Negros].

1927, Stettin, ent. Ztg 88: 311 (♂ ♀). Type-locs: Tappani, and Kanshizei, Formosa.

collaris de Meijere. — Borneo, Philippines [Mindanao]; Australia, Guadalcanal, New Britain, New Guinea.

1908, Tijdschr. Ent. 51: 258 (♂ ♀). Type-locs: Seleo, Berlinsfalen, New Guinea; Hapotheron, I. Deslac, and Kinigunang, New Britain.

erythropyga Walker. — Moluccas.

1864, J. Linn. Soc. Lond. 7: 211 (♂). Type-loc: Mysol, Moluccas.

flava Sack. — Philippines [Luzon].

1926, Philipp. J. Sci. 29: 582 (♂). Type-loc: Manila, Luzon, Philippines.

formosana Shiraki. — Formosa.

1930, Mem. Fac. Agric. Taihoku imp. Univ. 1: 144 (♂ ♀). Type-locs: Taiheisan and Tenchosan, Formosa.

inficita Walker. — Celebes.

1861, J. Linn. Soc. Lond. 5: 303 (♂). Type-loc: Tidon, Celebes.

javana Macquart. — Java.

1842, Dipt. exot. 2(2): 32 (92) (♂). Type-loc: Java.

marginata de Meijere. — Simeuluë Island.

1915, Tijdschr. Ent. 58 (Suppl.); 35 (♂ ♀). Type-locs: Pulu Babi, and Sinabang, Simalur.

meijerei van der Goot. — Simeuluë Island.

1964, Beaufortia 10: 219 (n. name for *chalybaeus* de Meijere).

chalybaeus de Meijere, 1915, Tijdschr. Ent. 58 (Suppl.): 30 (♀; preocc. Macquart 1842). Type-loc: Sinabang, Simalur.

neptunus de Meijere. — Java.

1911, Tijdschr. Ent. 54: 340 (♂). Type-loc: Muara Antjol nr Batavia (Djakarta), Java.

penangensis Curran. — Malaya.

1931, J. fed. Malay St. Mus. 16: 337 (♀). Type-loc: Penang Island, Malaya.

perakensis Curran. — Malaya.

1928, J. fed. Malay St. Mus. 14: 306 (♀). Type-loc: Jor Camp (2000 ft), Perak, Malaya.

polychromata Brunetti. — India [W Bengal].

1923, Fauna Br. India, Dipt. 3: 180 (♂ ♀). Type-loc: Calcutta, India.

resoluta Walker. — Sula Islands; Australia, Guadalcanal, Moluccas (Aru, Kai), New Guinea.

1858, J. Linn. Soc. Lond. 3: 95 (♂ ♀). Type-loc: Aru Islands, Moluccas.

sacki van der Goot. — Formosa.

1964, Beaufortia 10: 220 (n. name for *semisplendens* Shiraki).

semisplendens Shiraki, 1930, Mem. Fac. Agric. Taihoku imp. Univ. 1: 140 (♂ ♀; preocc. Sack 1926). Type-locs: Kotosh, Kankau, Sokutsu, and Kosempo, Formosa.

semicirculus Walker. — Not Oriental (see van der Wulp, 1899, Tijdschr. Ent. 42: 52); Neotropical.

simulata Brunetti. — India [Madras].

1923, Fauna Br. India, Dipt. 3: 177 (♂ ♀). Type-loc: Ootacamund, India.

tomentosa Macquart. — Java.

1848, Dipt. exot. Suppl. 3: 39 (199) (♀). Type-loc: Java.

violacea Macquart. — Java.

1848, Dipt. exot. Suppl. 3: 40 (200) (♀). Type-loc: Java.

Genus HELOPHILUS Meigen

Helophilus Meigen, 1822, Syst. Beschr. europ. zweifl. Insekten. 3: 368. Type-species, *Musca pendula* Linnaeus, by designation of Curtis (1832: pl. 429).

notabilis Macquart. — Java.

1842, Dipt. exot. 2(2): 63 (123) (♀). Type-loc: Java.

Unplaced species of *Helophilus* Meigen

aeneus Brunetti. — Nepal.

1907, Rec. Indian Mus. 1: pl. xii, fig. 15 (♀; as *aenous*; 1908, Rec. Indian Mus. 2: 66 (descr.); 1923, Fauna Br. India, Dipt. 3: 206 (emend.)). Type-loc: Soondrijal, Nepal.

aenous, error.

celeber Osten Sacken. — Philippines.

1882, Berl. ent. Z. 26: 116 (♂). Type-loc: Philippines.

conclusus Walker. — Celebes.

1860, J. Linn. Soc. Lond. 4: 119 (♂). Type-loc: Makasar, Celebes.

consors Walker. — Celebes.

1860, J. Linn. Soc. Lond. 4: 119 (♂). Type-loc: Makasar, Celebes.

Genus KEDA Curran

Keda Curran, 1931, J. fed. Malay St. Mus. 16: 331. Type-species, *Eristalis simpliciceps* de Meijere, by original designation, = *orientalis* (Rondani).

orientalis (Rondani). — Borneo, Java, Malaya; New Guinea.

1875, Annali Mus. civ. Stor. nat. Giacomo Doria 7: 421 (♀; *Eristalomyia*).

Type-loc: Sarawak, Borneo. N. comb.

simpliciceps de Meijere, 1914, Tijdschr. Ent. 57: 146 (♀; *Eristalis*). Type-loc: Batavia (Djakarta), Java. N. synonym.

Genus KERTESZIOMYIA Shiraki

Kertesziomyia Shiraki, 1930, Mem. Fac. Agric. Taihoku imp. Univ. 1: 151. Type-species, *Eristalis violascens* Kertész, by original designation.

fulgens (Macquart). — Java; Australia.

1850, Dipt. exot. Suppl. 4: 135 (439) (♂; *Eristalis*). Type-loc: Java.

violascens (Kertész). — Formosa.

1913, Annls hist.-nat. Mus. natn. hung. 11: 282 (♂ ♀; *Eristalis*). Type-locs: Kosempo, Fushoshio, Polisha, and Alikang, Formosa.

Genus MALLOTA Meigen

Mallota Meigen, 1822, Syst. Beschr. europ. zweifl. Insekts. 3: 377. Type-species, *Syrphus fuciformis* Fabricius, by designation of Rondani (1845:452).

Imatisma Macquart, 1842, Dipt. exot. 2(2): 67 (127). Type-species, *Eristalis posticatus* Fabricius, by original designation.

Tigridemyia Bigot, 1882, Annls Soc. ent. Fr. (6) 2 (Bull.): cxxi. Type-species, *pictipes* Bigot, by monotypy, = *curvigator* (Macquart). N. synonym.

Trigridiamyia, emendation.

Tigridomyia, emendation.

Teuchomerus Sack, 1922, Arch. Naturgesch. (A) 87(11): 266. Type-species, *Polydonta orientalis* Brunetti, by original designation, = *curvigator* (Macquart). N. synonym.

Helophilus, subg. *Paramesembrius* Shiraki, 1930, Mem. Fac. Agric. Taihoku imp. Univ. 1: 176. Type-species, *Tubifera abdominalis* Sack, by designation of Shiraki (1969: 227). N. synonym.

Mallota, subg. *Paramallota* Shiraki, 1930, Mem. Fac. Agric. Taihoku imp. Univ. 1: 191. Type-species, *haemorrhoidalis* Sack, by original designation.

Klossia Curran, 1931, J. fed. Malay St. Mus. 16: 370. Type-species, *dimidiata* Curran, by original designation, = *singularis* (Walker). N. synonym.

abdominalis (Sack). — Formosa; Japan.

1927, Stettin. ent. Ztg 88: 314 (♂; *Tubifera*). Type-locs: Suisharyo, and Kankau, Formosa. N. comb.

curvigator (Macquart). — Borneo, Ceylon, China, Formosa, India, Malaya.

1842, Dipt. exot. 2(2): 62 (122) (♂; *Helophilus*). Type-loc: Java. N. comb.
interveniens Walker, 1860, J. Linn. Soc. Lond. 4: 120 (♂; *Merodon*). Type-loc: Makasar, Celebes.

pictipes Bigot, 1882, Annls Soc. ent. Fr. (6) 2 (Bull.): cxxi (♂; *Tigridemyia*). Type-loc: Java.

orientalis Brunetti, 1908, Rec. Indian Mus. 2: 74 (♂; *Polydonta*). Type-loc: "Inde."

formosana Shiraki. — Formosa.

1930, Mem. Fac. Agric. Taihoku imp. Univ. 1: 196 (♂ ♀). Type-locs: Horisha, Koshun, and Kosempo, Formosa.

fulvula Curran. — Malaya.

1928, J. fed. Malay St. Mus. 14: 288 (♀). Type-loc: Sungai Bertam (4800 ft), Pahang, Malaya.

haemorrhoidalis Sack. — Formosa.

1927, Stettin. ent. Ztg 88: 317 (♂ ♀). Type-locs: Kosempo, and Chosokai, Formosa.

horishana Shiraki. — Formosa.

1930, Mem. Fac. Agric. Taihoku imp. Univ. 1: 194 (♂ ♀). Type-locs: Horisha, Toa Tsui Kutsu, Fuhosho, Kosempo, and Suisharyo, Formosa.

hysopia Vockeroth (for *horishanus* Shiraki). — Formosa. N. name.

horishanus Shiraki, 1930, Mem. Fac. Agric. Taihoku imp. Univ. 1: 178 (♂ ♀; *Helophilus*; preocc. Shiraki 1930). Type-loc: Horisha, Formosa. N. comb.

maculata Curran. — Thailand.

1928, J. fed. Malay St. Mus. 14: 289 (♀). Type-loc: Khao Luang (5800 ft), Nakhon Si Thammarat, Thailand.

matolla Knutson (for *orientalis* Macquart). — Formosa, Malaya. N. name.

orientalis Macquart, 1842, Dipt. exot. 2(2): 69 (129) (♂; *Imatisma*; preocc. Wiedemann 1824). Type-loc: "Indes Orientales."

nebulipennis (de Meijere). — Java.

1914, Tijdschr. Ent. 57: 145 (♀; *Eristalis*). Type-loc: Gunung Gede, Java. N. comb.

orientalis (Wiedemann). — Formosa, India [Sikkim, W Bengal], Java, Laos, Malaya.

1824, Anlecta Ent.: 38 (♂; *Eristalis*). Type-loc: Java.

quadrangulum (de Meijere). — Malaya.

1915, Tijdschr. Ent. 58 (Suppl.): 33 (♀; *Eristalis*). Type-loc: Sinabang, Malaya. N. comb.

rufipes Brunetti. — India [W Bengal].

1913, Rec. Indian Mus. 9: 271 (♂). Type-loc: Singla, Darjeeling District, India.

singularis (Walker). — Borneo, Malaya.

1857, J. Linn. Soc. Lond. 1: 17 (♀; *Eristalis*). Type-loc: Singapore, Malaya. N. comb.

dimidiata Curran, 1931, J. fed. Malay St. Mus. 16: 370 (♀; *Klossia*). Type-loc: Bettutan, nr Sandakan, N Borneo.

sufficiens Curran. — Malaya.

1928, J. fed. Malay St. Mus. 14: 291 (♀). Type-loc: Lubok Tamang (3500 ft), Pahang, Malaya.

vilis (Wiedemann). — Ceylon, India, Java, Thailand.

1830, Aussereurop. zweifl. Insekts. 2: 164 (♂; *Eristalis*). Type-loc: Java. N. comb.

eristaloides Curran, 1928, J. fed. Malay St. Mus. 14: 293 (♂; preocc. Loew 1856). Type-loc: Khao Luang (5800 ft), Nakhon Si Thammarat, Thailand. N. synonym.

malayana Curran, 1931, J. fed. Malay St. Mus. 16: 328 (n. name for *eristaloides* Curran).

Genus MESEMBRIUS Rondani

- Mesembrius* Rondani, 1857, Dipterol. ital. Prodr. 2: 50. Type-species, *Helophilus peregrinus* Loew, by monotypy.
- Eumerosyrphus* Bigot, 1882, Annls Soc. ent. Fr. (6) 2 (Bull.): cxxviii. Type-species, *indianus* Bigot, by monotypy, = *bengalensis* (Wiedemann).
- References: Curran 1925, 1947 (keys); Bhatia 1931 (biol., imm. stages).
atrolaneus Edwards. — Sumatra.
 1919, J. fed. Malay St. Mus. 8: 43 (♂). Type-loc: Sandaran Agong, Korinchi Lake, Sumatra.
- bengalensis* (Wiedemann). — SE Asia to New Guinea and northern Australia.
 1819, Zool. Mag. (Wied.) 1: 16 (♂ ♀; *Eristalis*). Type-loc: Bengal, India.
pilipes Doleschall, 1857, Natuurk. Tijdschr. Ned.-Indië 14: 410 (sex?; *Helophilus*). Type-loc: middle Java.
indianus Bigot, 1882, Annls Soc. ent. Fr. (6) 2 (Bull.): cxxviii (♂; *Eumerosyrphus*). Type-loc: India.
formosanus Shiraki. — Formosa.
 1930, Mem. Fac. Agric. Taihoku imp. Univ. 1: 172 (♂ ♀). Type-locs: Shinchiku, Horisha, Jukirin, Taihorin, Namakaban, Tabo, and Kotosho, Formosa.
- gigas* Curran. — Malaya.
 1928, J. fed. Malay St. Mus. 14: 296 (♀). Type-loc: Lubok Kedondong, NW of Mt Ophir (200 ft), Malacca-Johore border, Malaya.
- insignis* (Walker). — Borneo, Burma, Java, Malaya.
 1857 (March 1), J. Linn. Soc. Lond. 1: 17 (♀; *Helophilus*). Type-loc: Singapore, Malaya.
insignis Doleschall, 1857 (March 16), Natuurk Tijdschr. Ned.-Indië 14: 409 (sex?; *Helophilus*). Type-loc: Karangbollong, middle Java.
doleschalli Brunetti, 1908, Rec. Indian Mus. 2: 68 (n. name for *insignis* Doleschall).
- niveiceps* (de Meijere). — Formosa, Java, Ryukyu Islands.
 1908, Tijdschr. Ent. 51: 236 (♂; *Helophilus*). Type-loc: Java.
- quadrivittatus* (Wiedemann). — India, Java; Moluccas (Aru).
 1819, Zool. Mag. (Wied.) 1: 17 (♂; *Eristalis*). Type-loc: Tranquebar, Madras, India (Curran, 1930: 1).
- tuberosus* Curran. — China [Fukien].
 1925, Am. Mus. Novit. 200: 5 (♂ ♀). Type-loc: Yen-ping, Fukien, China.
- vestitus* (Wiedemann). — Java, Sumatra.
 1824, Analecta Ent.: 36 (♂ ♀; *Eristalis*). Type-loc: Java.
- wulpi van der Goot. — Celebes, Formosa, Java, Philippines [Luzon, Negros]; Moluccas (Aru), New Guinea.
 1964, Beaufortia 10: 220 (n. name for *Helophilus albiceps* van der Wulp).
albiceps van der Wulp, 1867, Tijdschr. Ent. 11: 116 (♂ ♀; *Helophilus*; preocc. Macquart 1846). Type-locs: Celebes; and Aru Islands, Moluccas.

Genus PHYTOMIA Guérin-Méneville

- Pachycephalus* Wiedemann, 1830, Aussereurop. zweifl. Insekts. 2: 152 (preocc. Stephens 1826). Type-species, *Eristalis chrysopygus* Wiedemann, by present designation.

Syphidae

- Phytomia* Guérin-Méneville, 1834, in Bélanger, Voyage Ind.-Orient. 7:509. Type-species, *Eristalis chrysopygus* Wiedemann, by monotypy.
- Megaspis* Macquart, 1842, Dipt. exot. 2(2): 27 (87). Type-species, *Eristalis chrysopygus* Wiedemann, by designation of Brunetti 1923: 196.
- References: Brunetti 1908; de Meijere 1908 (keys).
- Subgenus PHYTOMIA Guérin-Méneville
- aesymnus* (Walker). — Borneo, ?India, Java, Malaya, Sumatra.
 1849, List Dipt. Br. Mus. 3: 630 (♂; *Eristalis*). Type-loc: East Indies.
flavofasciatus Macquart, 1850, Dipt. exot. Suppl. 4: 136 (440) (♂ ♀; *Eristalis*). Type-loc: Java. N. synonym.
- ursina* Jaennicke, 1867, Abh. senckenb. naturf. Ges. 6: 401 (♀; *Eristalis*). Type-loc: Java.
- argyrocephala* (Macquart). — Burma, India [Madras, Mysore, W Bengal], Java, Laos, Sumatra, Sumbawa.
 1842, Dipt. exot. 2(2): 45 (105) (♂ ♀; *Eristalis*). Type-loc: "Indes orientales."
transversus Brunetti, 1908, Rec. Indian Mus. 2: 73 (♂ ♀; *Megaspis*). Type-locs: Calcutta, and Bangalore, India.
- chrysopyga* (Wiedemann). — Burma, India [Assam], Java, Laos, Malaya, Sumatra, Thailand.
 1819, Zool. Mag. (Wied.) 1: 15 (♂; *Eristalis*). Type-loc: Java.
- aurata* Macquart, 1834, Hist. nat. Ins. Dipt. 1: 494 (♂; *Volucella*). Type-loc: Java.
- errans* (Fabricius). — China, throughout SE Asia; Japan.
 1787, Mantissa Insect. 2: 337 (♀; *Syrphus*). Type-loc: China.
- varipes* Macquart, 1842, Dipt. exot. 2(2): 46 (106) (♂ ♀; *Eristalis*). Type-locs: "Indes orientales" and China.
- amphicrates* Walker, 1849, List Dipt. Colln Br. Mus. 3: 623 (♂ ♀; *Eristalis*). Type-locs: N Bengal; Java; East Indies; and China.
- plistoanax* Walker, 1849, List Dipt. Colln Br. Mus. 3: 628 (♂; *Eristalis*). Type-loc: Philippine Islands.
- ayrus* Walker, 1849, List Dipt. Colln Br. Mus. 3: 629 (♂; *Eristalis*). Type-loc: Philippine Islands.
- macquarti* Doleschall, 1856, Natuurk. Tijdschr. Ned.-Indië 10: 410 (sex?; *Eristalis*). Type-loc: Java.
- sculptata* (van der Wulp). — Timor.
 1868, Tijdschr. Ent. 11: 113 (♀; *Megaspis*). Type-loc: Timor.
- tenebrica* Edwards. — Sumatra.
 1919, J. fed. Malay St. Mus. 8: 42 (♂). Type-loc: Sungai Kumbang, Sumatra.
- zonata* (Fabricius). — entire SE Asia; Japan, Korea, Manchuria.
 1787, Mantissa Insect. 2: 337 (♀; *Syrphus*). Type-loc: China.
- zonalis*, error or emendation.
- sinensis* Gmelin, 1790, Syst. Nat. 5: 2872 (sex?; *Musca*; unjustified n. name for *zonata* Fabricius).
- lata* Macquart, 1842, Dipt. exot. 2(2): 35 (95) (♀; *Eristalis*). Type-loc: "Patrie inconnue."
- andraemon* Walker, 1849, List Dipt. Colln Br. Mus. 3: 627 (♀; *Eristalis*). Type-locs: Sylhet, E Pakistan and Sikkim, India.

- datamus* Walker, 1849, List Dipt. Colln Br. Mus. 3: 628 (♀; *Eristalis*). Type-loc: unknown.
- babytace* Walker, 1849, List Dipt. Colln Br. Mus 3: 629 (♂; *Eristalis*). Type-loc: Philippine Islands.
- exterus* Walker, 1852, Insecta Saundersiana 1: 248 (♀; *Eristalis*). Type-loc: East Indies.
- cingulata* Vollenhoven, 1863, Versl. Meded. K. Akad. wet. Amst. 15: 12 (sex?; *Megaspis*). Type-loc: Japan.

Subgenus DOLICHOMERUS Macquart

- Dolichomerus* Macquart, 1850, Dipt. exot. Suppl. 4: 131 (435). Type-species, *Syrphus crassus* Fabricius, by monotypy.
- crassa* (Fabricius). — Celebes, Ceylon, India [Andhra Pradesh, Assam, Madras], Laos, Malaya, Nepal, Thailand.
- 1787, Mantissa Insect. 2: 334 (sex?; *Syrphus*). Type-loc: Tranquebar, Madras, India.
- mégacephalus* Fabricius, 1798, Suppl. Ent. Syst.: 561 (sex?; *Syrphus*). Type-loc: "India oriental."

Genus PSEUDERISTALIS Shiraki

- Eristalis*, subg. *Pseuderistalis* Shiraki, 1930, Mem. Fac. Agric. Taihoku imp. Univ. 1: 147. Type-species, *bicolor* Shiraki, by original designation.
- bicolor* (Shiraki). — Formosa.
- 1930, Mem. Fac. Agric. Taihoku imp. Univ. 1: 148 (♂ ♀; *Eristalis*). Type-loc: Suisha, Formosa.
- fascipennis* Thompson (for *maculipennis* de Meijere). — Burma, India [Assam], Java, Philippines [Mindoro, Negros, Mindanao]. N. name.
- maculipennis* de Meijere, 1908, Tijdschr. Ent. 51: 261 (♂; *Eristalis*; preocc. Townsend 1897). Type-loc: Lawang, E Java.
- nigra* (Wiedemann). — throughout SE Asia; Moluccas (Mysol), New Guinea.
- 1824, Analecta Ent.: 38 (♀; *Eristalis*). Type-loc: Java.
- bombooides* Walker, 1860, J. Linn. Soc. Lond. 4: 119 (♂ ♀; *Eristalis*). Type-loc: Makasar, Celebes.
- obscurata* Walker, 1861, J. Linn. Soc. Lond. 5: 239 (♂ ♀; *Eristalis*). Type-loc: Dorey, New Guinea.
- tortuosa* Walker, 1861, J. Linn. Soc. Lond. 5: 266 (♂; *Eristalis*). Type-loc: Tond, Celebes.
- pura* Curran. — Malaya.
- 1928, J. fed. Malay. St. Mus. 14: 305 (♀). Type-loc: Ginting Sempak, Selangor-Pahang border, Malaya.
- semiplendens* (Sack). — Philippines [Luzon, Palawan, Leyte, Negros, Mindanao].
- 1926, Philipp. J. Sci. 29: 584 (♀; *Eristalis*). Type-loc: Cuernos Mts, Negros Occidental, Philippines. N. comb.
- velutina* (Sack). — Philippines [Negros].
- 1926, Philipp. J. Sci. 29: 586 (♀; *Eristalis*). Type-loc: Cuernos Mts, Negros Occidental, Philippines. N. comb.

Genus SENASPIST Macquart

- Senaspis* Macquart, 1850, Dipt. exot. Suppl. 4: 133 (437). Type-species, *flaviceps* Macquart, by monotypy.
- dentipes* (Macquart). — Not Oriental; Ethiopian.
- 1842, Dipt. exot. 2(2): 37 (97) (♂; *Eristalis*). Type-loc: "Java." Senior synonym of *aesculus* (Walker), a common Ethiopian species.

Tribe MILESINI

Genus BRACHYPALPUS Macquart

- Brachypalpus* Macquart, 1834, Hist. nat. Ins. Dipt. 1: 523. Type-species, *tuberculatus* Macquart, by designation of Rondani (1845: 456), = *valgus* (Panzer).
- dives* Brunetti. — India [Assam].
- 1908, Rec. Indian Mus. 2: 86 (♂). Type-loc: Kohima, Assam, India.
- trilineatus* Hull. — Java.
- 1943, J. Wash. Acad. Sci. 33: 43 (♀). Type-loc: Mt Gede, Tjibodas, Java.

Genus CHEIROXYLOTA Hull

- Xylota*, subg. *Cheiroxylota* Hull, 1949, Trans. zool. Soc. Lond. 26: 361. Type-species, *dimidiata* Brunetti, by original designation.
- dimidiata* (Brunetti). — Nepal, W Pakistan.
- 1923, Fauna Br. India, Dipt. 3: 232 (♀; *Xylota*). Type-loc: Dungagli, Hazara District, India.

Genus CRIORHINA Meigen

- Penthesilea* Meigen, 1800, Nouv. Class. Mouches: 35. Type-species, *Musca ruficauda* DeGeer, by subsequent monotypy (Bezzi, in Hendel 1908: 67); misidentification, = *ranunculi* (Panzer). Suppressed by I.C.Z.N. 1963, Opinion 678: 339.
- Criorrhina* Meigen, 1822, Syst. Beschr. europ. zweifl. Insek. 3: 236. Type-species, *Syrphus astilicus* Fallén, by designation of Westwood (1840: 136).
- Criorrhina*, emendation.
- bombooides* Hull. — India [Assam].
- 1944, Ann. Mag. nat. Hist. (11) 11: 37 (♂). Type-loc: Shillong, Assam, India.
- crioartctos* Hull. — Burma.
- 1944, Ann. Mag. nat. Hist. (11) 11: 36 (♀). Type-loc: Burma.
- cyaniventris* (Sack). — Philippines [Negros].
- 1926, Philipp. J. Sci. 29: 588 (♂; *Penthesilea*). Type-loc: Fabrica, Negros Occidental, Philippines.
- dentata* Brunetti. — India.
- 1908, Rec. Indian Mus. 2: 87 (♂). Type-loc: Kodiali, Simla District, India.
- excavata* Curran. — India [Uttar Pradesh].
- 1929, Ann. ent. Soc. Am. 22: 498 (♂). Type-loc: Deoban, Chakrata, India.

formosana Shiraki. — Formosa.

1930, Mem. Fac. Agric. Taihoku imp. Univ. 1: 51 (♀). Type-loc: Rantai-zan, Formosa.

imitator Brunetti. — India [Kashmir, Uttar Pradesh].

1915, Rec. Indian Mus. 11: 236 (♀). Type-loc: W Himalaya, Garhwal District, India.

interrupta Brunetti. — W Pakistan.

1923, Fauna Br. India, Dipt. 3: 287 (♂). Type-loc: Dungagali, Hazara District, (W Pakistan).

pallidipes Curran. — India [Uttar Pradesh].

1929, Ann. ent. Soc. Am. 22: 498 (♂). Type-loc: Deoban, Chakrata, India.

pallipilosa Hull. — India [Kashmir].

1944, Ann. Mag. nat. Hist. (11) 11: 34 (♂). Type-loc: Gulmarg, Kashmir, India.

ssp. *bicolorata* Hull. — India [Kashmir].

1950, Ann. Mag. nat. Hist. (12) 3: 610 (♀). Type-loc: Gulmarg, Kashmir, India.

rubripes (Matsumura). — Formosa; Japan.

1916, Thousand Insects Japan 2: 218 (♀; *Mallota*). Type-loc: Kyoto, Honshu, Japan.

rubropilosa Hull. — India [Kashmir].

1950, Ann. Mag. nat. Hist. (12) 3: 608 (♀). Type-loc: Gulmarg, Kashmir, India.

setosa Shiraki. — Formosa.

1930, Mem. Fac. Agric. Taihoku imp. Univ. 1: 54 (♀). Type-loc: Karenko, Formosa.

shirakii van der Goot. — Formosa.

1964, Beaufortia 10: 220 (n. name for *apiformis* Shiraki).

apiformis Shiraki, 1930, Mem. Fac. Agric. Taihoku imp. Univ. 1: 52 (♀; preocc. Macquart 1829). Type-loc: Karenko, Formosa.

simiodes (Brunetti). — India.

1908, Rec. Indian Mus. 2: 83 (♂; *Deineches*). Type-loc: "Inde."

spinitarsis Curran. — India [Uttar Pradesh].

1929, Ann. ent. Soc. Am. 22: 499 (♂). Type-loc: Rishikesh, Dehra Dun, India.

trifasciata Shiraki. — Formosa.

1930, Mem. Fac. Agric. Taihoku imp. Univ. 1: 56 (♂). Type-loc: Rantai-zan, Formosa.

tripilosa Coe. — India [Uttar Pradesh].

1964, Bull. Br. Mus. nat. Hist. (Ent.) 15: 279 (♀). Type-loc: Barasu, Kumaon, India.

vivida Brunetti. — India [Kashmir].

1923, Fauna Br. India, Dipt. 3: 285 (♀). Type-loc: Kashmir, India.

Genus KORINCHIA Edwards

Korinchia Edwards, 1919, J. fed. Malay St. 8: 39. Type-species, *klossi* Edwards, by designation of Edwards, in Brunetti (1923: 224).

Palumbia (4)

apicalis Shiraki. — Formosa.

1930, Mem. Fac. Agric. Taihoku imp. Univ. 1: 157 (♂). Type-loc: Karenko, Formosa.

formosana Shiraki. — Formosa.

1930, Mem. Fac. Agric. Taihoku imp. Univ. 1: 154 (♂ ♀). Type-locs: Shinchiku, Arisan, Karenko, and Tabani, Formosa.

klossi Edwards. — Sumatra.

1919, J. fed. Malay St. Mus. 8: 40 (♂). Type-loc: Sungai Kumbang, Sumatra.

minor (de Meijere). — Sumatra.

1919, Bijdr. Dierk. 21: 27 (♂; *Milesia*). Type-loc: Peak of Korinchi, Sumatra.

monticola Edwards. 1919, J. fed. Malay St. Mus. 8: 41 (♂). Type-loc: Korinchi Peak, Sumatra.

nova Hull. — China.

1937, J. Wash. Acad. Sci. 27: 175 (♀). Type-loc: Ningyuensu, China.

pendleburyi Curran. — Borneo.

1931, J. fed. Malay St. Mus. 16: 373 (♂). Type-loc: Mt Kinabalu, Kamborangah (7200 ft), N Borneo.

robinsoni Edwards. — Sumatra.

1919, J. fed. Malay St. Mus. 8: 41 (♀). Type-loc: Sungai Kumbang, Sumatra.

rufa Hervé-Bazin. — India [Madras].

1922, Bull. Soc. ent. Fr. 1922: 122 (♂ ♀). Type-loc: Kodaikanal, S India.

simulans (de Meijere). — Java, Laos, Sumatra.

1914, Tijdschr. Ent. 57: 144 (♂ ♀; *Milesia*). Type-locs: Gunung Gede, and Gunung Ungaran, Java.

aurata Hervé-Bazin, 1922, Bull. Soc. ent. Fr. 1922: 213 (♀). Type-loc: Mt Gede, Java.

Genus LYCASTRIS Walker

Lycastris Walker, 1857, Trans. ent. Soc. Lond. (n. Ser.) 4: 155. Type-species *albipes* Walker, by monotypy.

Xiphopheromyia Bigot, 1892, Wien. Ent. Ztg 11: 161. Type-species, *glossata* Bigot, by monotypy, = *albipes* Walker.

Reference: Coe 1964 (key).

albipes Walker. — India.

1857, Trans. ent. Soc. Lond. (n. ser.) 4: 155 (♀). Type-loc: "Hindostan."

glossata Bigot, 1892, Wien. ent. Ztg 11: 161 (♂; *Xiphopheromyia*). Type-loc: Sabateo, India.

austeni Brunetti. — India [W Bengal].

1923, Fauna Br. India, Dipt. 3: 279 (♂ ♀). Type-loc: Darjeeling, India.

cornuta Enderlein. — Formosa.

1911, Stettin ent. Ztg 72: 236 (♀). Type-loc: Formosa.

flavohirta Brunetti. — India [W Bengal].

1907, Rec. Indian Mus. 1: pl. xi, figs. 16 & 17; 1908, Rec. Indian Mus. 2: 85 (♂). Type-loc: Darjeeling, India.

griseipennis Cœ. — India [Assam].

1964, Bull. Br. Mus. nat. Hist. (Ent.) 15: 281 (♂). Type-loc: Mishmi Hills, Delai Valley, Cha Che, Assam, NE India.

Genus MILESIA Latreille

Milesia Latreille, 1804, Nouv. Dict. Hist. nat. 24 (Tab. méth. Ins.): 194. Type-species, *Syrphus crabroniformis* Fabricius, by designation of Williston (1887: 254).

To preserve the long and almost universal usage of *Milesia* in the sense of *Syrphus crabroniformis* Fabricius, as type-species, the International Commission on Zoological Nomenclature should be asked to suspend the rules and suppress Rondani's (1844, Nuovi Annali Sci. nat. Bologna (2) 2: 455) designation of *Musca diophthalma* Linnaeus (type-species of *Spilomyia* Meigen) and to place *Milesia* Latreille on the Official list of Generic Names with *Syrphus crabroniformis* Fabricius as type-species (see Williston 1887: 254–255).

ammochrysus Séguy. — Viet Nam.

1948, Notes ent. chin. 12: 166 (♂). Type-loc: Caleu, Annam (N Viet Nam).

apicalis Snellen van Vollenhoven. — Java.

1863, Versl. Meded. K. Akad. wet. Amst. 15: 13 (♂). Type-loc: Java.

apsycta Séguy. — China [Fukien].

1948, Notes ent. chin. 12: 167 (♂; as *apsyctos*). Type-loc: "Chine: Fou-kién, San-tao-ho, Ile des Araignées."

This name is a Greek adjective; in accordance with Article 11b of the International Code we have latinized it.

balteata Kertész. — India [Assam, Sikkim], Laos, Malaya, Thailand.

1901, Természetr. Füz. 24: 414 (♂ ♀). Type-loc: Sikkim, India.

himalayensis Brunetti, 1908, Rec. Indian Mus. 2: 82 (♂ ♀). Type-locs: Assam and Sikkim, India.

bigoti Osten Sacken. — Philippines [Luzon].

1882, Berl. ent. Z. 26: 190 (♂ ♀). Type-loc: Philippines.

brunettii Hervé-Bazin. — Laos, Nepal.

1923, Bull. Soc. ent. Fr. 1923: 26 (♀). Type-locs: Laos; and Himalaya.

brunnea Hervé-Bazin. — Laos.

1923, Bull. Soc. ent. Fr. 1923: 26 (♂). Type-loc: Laos.

callida Curran. — Borneo, Malaya.

1928, J. fed. Malay St. Mus. 14: 181 (♂). Type-loc: Payang, 'beluka' (500 ft), Singapore, Malaya.

conspicienda Walker. — Celebes, Philippines [Luzon, Sibuyan, Mindanao].

1860, J. Linn. Soc. Lond. 4: 118 (♂ ♀). Type-loc: Makasar, Celebes.

conspicua Curran. — Malaya.

1928, J. fed. Malay St. Mus. 14: 178 (♂). Type-locs: Gunong Tahan, Tangga Duablas (5700 ft) and Cameron Highlands, Rhododendron Hill (5200 ft), Pahang, Malaya.

decora Brunetti. — India [Assam].

1923, Fauna Br. India, Dipt. 3: 274 (♀). Type-loc: Lower Ranges, N Khasi Hills, Assam, India.

diardi Snellen van Vollenhoven. — Borneo.

1863, Versl. Meded. K. Akad. wet. Amst. 15: 14 (♂). Type-loc: Borneo.

doriae (Rondani). — Borneo.

1875, Annali Mus. civ. Stor. nat. Giacomo Doria 7: 422 (♂; *Sphyxea*). Type-loc: Sarawak, Borneo.

excelda Curran. — Malaya, Thailand.

1928, J. fed. Malay St. Mus. 14: 184 (♂). Type-locs: Nakhon Si Thammarat, Khao Ram (1200–1300 ft), Thailand; and Selangor, Kuala Lumpur, Gombak Valley, Malaya.

ferruginea Brunetti. — India [Assam, Sikkim, Uttar Pradesh, W Bengal], Laos.

1913, Rec. Indian Mus. 9: 268 (♀). Type-loc: Darjeeling District, India.

fissipennis Speiser. — Formosa; Japan.

1911, Jb. nassau Ver. Naturk. 64: 241 (♂). Type-loc: Chip-Chip, Formosa.

fissiformis, error.

flavifacies (Bigot). — Borneo.

1875, Annls Soc. ent. Fr. (5) 5: 471 (♂; *Sphyxea*). Type-loc: Sarawak, Borneo.

fulvipes (Bigot). — Java.

1883, Annls Soc. ent. Fr. (6) 3: 340 (♂ ♀; *Sphyxea*). Type-loc: Java.

fuscicosta (Bigot). — Borneo.

1875, Annls Soc. ent. Fr. (5) 5: 469 (♂ ♀; *Sphyxea*). Type-loc: Sarawak, Borneo.

insistens Curran. — Borneo.

1931, J. fed. Malay St. Mus. 16: 345 (♂ ♀). Type-loc: Mt Kinabalu, Lumu Lumu (5500 ft), N Borneo.

limbipennis Macquart. — Java.

1848, Dipt. exot. Suppl. 3: 42 (202) (♀). Type-loc: Java.

macularis Wiedemann. — Borneo, Java, Malaya, Sumatra.

1824, Analecta Ent.: 34 (♂). Type-loc: Java.

metallica Curran. — Borneo.

1931, J. fed. Malay St. Mus. 16: 347 (♂). Type-loc: Bettutan, nr Sandakan, N Borneo.

meyeri Jaennicke. — Java.

1867, Abh. senckenb. naturf. Ges. 6: 403 (♂). Type-loc: Java.

pendleburyi Curran. — Malaya.

1928, J. fed. Malay St. Mus. 14: 179 (♂). Type-loc: Jor Camp (1500 ft), Batang Padang, Perak, Malaya.

reinwardtii Wiedemann. — Borneo, Java, Malaya, Philippines [Luzon].

1824, Analecta Ent.: 33 (♂). Type-loc: Java.

ritsemae Osten Sacken. — Philippines [Luzon].

1882, Berl. ent. Z. 26: 191 (♀). Type-loc: Philippines.

semifulva de Meijere. — Burma, India [Assam, W Bengal], Java, Malaya, Sumatra.

1904, Bijdr. Dierk. 17: 99 (♂). Type-loc: Darjeeling, India.

gigas Guérin-Méneville, 1834, in Bélanger, Voyage Ind. Orient: 508 (♂; preocc. Rossi 1790). Type-loc: "côte du Coromandel" (India).

gigas Macquart, 1834, Hist. nat. Ins. Dipt. 1: 533 (♂). Type-loc: Java.

semperi Osten Sacken. — Philippines [Luzon, Negros, Mindanao].

1882, Berl. Ent. Z. 26: 189 (♂ ♀). Type-loc: Philippines.

sexmaculata Brunetti. — India [Madras].

1915, Rec. Indian Mus. 11: 248 (♂). Type-loc: Trivandrum, Travancore State, India.

simillima Hull. — Java, Sumbawa.

1950, Ann. Mag. nat. Hist. (12) 3: 606 (♂ ♀). Type-loc: Sockaboemi, Java.

sinensis Curran. — China [Fukien].

1925, Am. Mus. Novit. 200: 3 (♂ ♀). Type-loc: Yen-ping, China.

tenuiformis Curran. — Anambas Islands, Borneo, Malaya.

1928, J. fed. Malay St. Mus. 14: 176 (♂ ♀). Type-locs: Kuala Lumpur, Selangor and Kuala Tahan (300 ft), Pahang, Malaya.

variegata Brunetti. — India [Assam, Sikkim], Laos.

1908, Rec. Indian Mus. 2: 80 (♂). Type-loc: Sikkim, India.

verticalis Brunetti. — India [Assam].

1923, Fauna Br. India, Dipt. 3: 269 (♂ ♀). Type-loc: Tura, Garo Hills, Assam, India.

vesparia Shiraki. — Formosa.

1930, Mem. Fac. Agric. Taihoku imp. Univ. 1: 118 (♂ ♀). Type-locs: Shinchiku, Horisha, Rono, Karenko, Fuhosho, and Kosempo, Formosa.

vespoides Walker. — Malaya.

1857, J. Linn. Soc. Lond. 1: 18 (♂). Type-loc: Singapore, Malaya.

zamiel Walker. — Borneo.

1857, J. Linn. Soc. Lond. 1: 123 (♀). Type-loc: Sarawak, Borneo.

Genus NEPENTHOSYRPHUS de Meijere

Nepenthosyrphus de Meijere, 1932, Tijdschr. Ent. 75 (Suppl.): 155. Type-species, *tobaicus* de Meijere, by original designation.

capitatus (Sack). — Sumatra.

1931, Arch. Hydrobiol. (Suppl.-Bd.) 8: 590 (♂ ♀; *Syritta*). Type-locs: "Heide von Huta Gindjang and Balige, Tobagebiet"; Sibolga, Sumatra.

oudemansi de Meijere. — Borneo.

1932, Tijdschr. Ent. 75 (Suppl.): 157 (♂ ♀). Type-loc: Pontianak, W Borneo.

tobaicus de Meijere. — Sumatra.

1932, Tijdschr. Ent. 75 (Suppl.): 159 (♂ ♀). Type-locs: Tinggi Radja and Haranggaol, Toba Sea, W Sumatra.

Genus SPILOMYIA Meigen

Spilomyia Meigen, 1803, Magazin InsektKde 2: 273. Type-species, *Musca diophthalma* Linnaeus (as *Syrphus diophthalmus* Fabricius), by designation of Coquillett (1910: 607).

chinensis Hull. — China [Yunnan].

1950, Ann. Mag. nat. Hist. (12) 3: 603 (♀). Type-loc: Mekong River, Yei-chih, Yunnan, SW China.

Genus SYRITTA Lepeletier and Serville

Xylota, subg. *Syritta* Lepeletier and Serville, in Latreille, et al., 1828, Encycl. méth. (Ins.) 10(2): 808. Type-species, *Musca pipiens* Linnaeus (as *Xylota pipiens* Meigen), by monotypy.

Reference: Hodson 1931 (biol., imm. stages).

hackeri Klocker. — Java; Australia.

1924, Mem. Qd Mus. 8: 59 (♂). Type-loc: Brisbane, Queensland, Australia.

indica (Wiedemann). — Formosa, India [Madras, W Bengal], Nepal.

1824, Analecta Ent. 33 (♂ ♀; *Eumerus*). Type-loc: East Indies.

rufifacies Bigot, 1883, Annls Soc. ent. Fr. (6) 3: 538 (♂). Type-loc: Pondicherry, India. N. synonym.

semorata Sack, 1913, Ent. Mitt. 2: 8 (♂). Type-loc: Tainan, Formosa. N. synonym.

maritima Hull. — Christmas Island.

1944, Ann. Mag. nat. Hist. (11) 11: 60 (♀). Type-loc: Christmas Island, Indian Ocean (Indonesia).

orientalis Macquart. — Ceylon, India [Madras], ?Java, ?Celebes; ?Amboina.

1842, Dipt. exot. 2(2): 76 (136) (♂). Type-loc: Pondicherry, Madras, India.

?*lutescens* Doleschall, 1856, Natuurk. Tijdschr. Ned.-Indië 10: 410 (sex?; *Senogaster*). Type-loc: Djokjokarta, Java.

?*ambainensis* Doleschall, 1858, Natuurk. Tijdschr. Ned.-Indië 17: 97 (sex?). Type-loc: Amboina, Moluccas.

?*illucida* Walker, 1860, J. Linn. Soc. Lond. 4: 121 (♀). Type-loc: Makasar, Celebes.

pipiens (Linnaeus). — Nepal; Holarctic.

1758, Syst. Nat. Ed. 10: 594 (sex?; *Musca*). Type-loc: Europe.

triangulifera Keiser. — Ceylon.

1958, Revue suisse Zool. 65: 226 (♂ ♀). Type-loc: Belihul Oya, Ceylon.

Genus TAKAOMYIA Hervé-Bazin

Takaomyia Hervé-Bazin, 1915, Annls Soc. ent. Fr. 83: 412. Type-species, *johannis* Hervé-Bazin, by monotypy.

formosana Shiraki. — Formosa.

1930, Mem. Fac. Agric. Taihoku imp. Univ. 1: 125 (♂ ♀). Type-loc: Horisha, Musha, Formosa.

johannis Hervé-Bazin. — Formosa, Thailand; Japan.

1915, Annls Soc. ent. Fr. 83: 413 (♂). Type-loc: "Mont Takao, près Hachioji," Japan.

Genus TEMNOSTOMA Lepeletier and Serville

Tritonia Meigen, 1800, Nouv. Class. Mouches: 33 (preocc. Cuvier 1798). Type-species, *Musca vespiformis* Linnaeus, by designation of Coquillett (1910: 617). Suppressed by I.C.Z.N., 1963, Opinion 678: 339.

Temnostoma Lepeletier and Serville, in Latreille, et al., 1828, Encycl. méth. (Ins.) 10: 518. Type-species, *Milesia bombylanis* Fabricius, by designation of Coquillett (1910: 612).

nigrimanus Brunetti. — India [Uttar Pradesh].

1915, Rec. Indian Mus. 11: 246 (♂). Type-loc: Garhwal District, India.

taiwanum Shiraki. — Formosa.
1930, Mem. Fac. Agric. Taihoku imp. Univ. 1: 104 (♀). Type-loc: Asahi, Formosa.

Genus XYLOTA Meigen

- Zelima* Meigen, 1800, Nouv. Class. Mouches: 34. Type-species, *Musca segnis* Linnaeus, by designation of Coquillett (1910: 621). Suppressed by I.C.Z.N., 1963, Opinion 678: 339.
- Helophilus* Meigen, 1803, Magazin Insektskde 2: 273. Type-species, *Musca sylvarum* Linnaeus (as *Syrphus sylvarum* Fabricius), by monotypy.
- Eumeros* Meigen, 1803, Magazin Insektskde 2: 273. Type-species, *Musca segnis* Linnaeus, by designation of Coquillett (1910: 541).
- Xylota* Meigen, 1822, Syst. Beschr. europ. zweifl. Insekts. 3: 211 (unjustified n. name for *Helophilus* Meigen). Type-species, *Musca sylvarum* Linnaeus, automatic.
- aeneimaculata* de Meijere. — Formosa; New Guinea.
1908, Tijdschr. Ent. 51: 227 (♂). Type-loc: Moroka, New Guinea.
- abosa* Séguay. — Laos.
1948, Notes ent. chin. 12: 164 (♀). Type-loc: Nam-mat, Laos.
- acoetes* Séguay. — China.
1948, Notes ent. chin. 12: 165 (♂). Type-locs: Che-mo, and Houa-chau, China.
- aethusa* Walker. — ?Java.
1849, List. Dipt. Colln Br. Mus. 3: 559 (♀). Type-loc: East Indies.
- annulata* Brunetti. — India [W Bengal], Laos, Malaya.
1913, Rec. Indian Mus. 9: 270 (♂ ♀). Type-locs: Singla, and Darjeeling, India.
- var. *ornatipes* (Sack). — Formosa.
1927, Stettin. Ent. Ztg 88: 317 (♂; *Zelima*). Type-loc: Toa Tsui Kutsu, Formosa.
- annulipes* de Meijere. — Sumatra.
1924, Tijdschr. Ent. 67 (Suppl.): 19 (♀). Type-loc: Tandj. Andalas, Sumatra.
- armipes* (Sack). — Formosa.
1922, Arch. Naturgesch. (A) 87(11): 269 (♂; *Zelima*). Type-locs: Sokutsu, and Fuhosho, Formosa.
- assamensis* Brunetti. — India [Assam].
1908, Rec. Indian Mus. 2: 79 (♀). Type-loc: Kohima, Assam, India.
- auronitens* Brunetti. — India [Assam].
1908, Rec. Indian Mus. 2: 78 (♂). Type-loc: Margherita, Assam, India.
- bimaculata* (Shiraki). — Formosa.
1930, Mem. Fac. Agric. Taihoku imp. Univ. 1: 85 (♂; *Zelima*). Type-loc: Taikan, Formosa.
- bistriata* Brunetti. — India [Mysore], Thailand, Viet Nam.
1915, Rec. Indian Mus. 11: 235 (♂ ♀). Type-loc: Parambikulam, India.
- brachypalpoides* (Shiraki). — Formosa.
1930, Mem. Fac. Agric. Taihoku imp. Univ. 1: 70 (♂; *Zelima*). Type-loc: Sinsuie, Formosa.

- brunettii* Curran. — Malaya.
1928, J. fed. Malay St. Mus. 14: 277 (♀). Type-loc: Lubok Tamang (3500 ft), Pahang, Malaya.
- calopus* Bigot. — Java.
1883, Annls Soc. ent. Fr. (6) 3: 543 (♂). Type-loc: Java.
- carbonaria* Brunetti. — India [Madras].
1923, Fauna Br. India, Dipt. 3: 240 (♂). Type-loc: Shevaroys, Yercaud, India.
- conformis* Walker. — Malaya, Philippines [Luzon, Mindanao].
1857, J. Linn. Soc. Lond. 1: 18 (♀). Type-loc: Singapore.
- coquilletti* Hervé-Bazin. — Formosa; Japan.
1915, Annls Soc. ent. Fr. 83: 409 (n. name for *cuprina* Coquillett).
cuprina Coquillett, 1898, Proc. U.S. natn. Mus. 21: 327 (♂; preocc. Bigot 1885). Type-loc: Japan.
- cupreiventris* Brunetti. — India [W Bengal], Laos.
1923, Fauna Br. India, Dipt. 3: 233 (♀). Type-loc: Surcil, Darjeeling District, India.
- cuprina* Bigot. — India.
1885, Annls Soc. ent. Fr. (6) 5: 247 (♂). Type-loc: "Indes."
- decora* de Meijere. — Java.
1914, Tijdschr. Ent. 57: 142 (♀). Type-loc: Gungung Gede, Java.
- doris* Curran. — Malaya.
1928, J. fed. Malay St. Mus. 14: 281 (♀). Type-loc: Gunung Berumban, (4000–4500 ft), Cameron Highlands, Pahang, Malaya.
- flavipes* (Sack). — Formosa.
1927, Stettin. Ent. Ztg 88: 316 (♀; *Zelima*). Type-loc: Toa Tsui Kutsu, Formosa.
- fo. Hull. — China [Yunnan].
1944, Proc. ent. Soc. Wash. 46(2): 45 (♂). Type-loc: San-Nen-Kai, Yunnan, SE China.
- formosana* Matsumura. — Formosa.
1916, Thousand Insects Japan 2: 251 (♂). Type-loc: Horisha, Formosa.
- makiana* (Shiraki). — Formosa, Nepal.
1930, Mem. Fac. Agric. Taihoku imp. Univ. 1: 65 (♂ ♀; *Zelima*). Type-locs: Arisan, Kosempo, Toyenmongai, and Fuhosho, Formosa.
- morna* Curran. — Borneo.
1931, J. fed. Malay St. Mus. 16: 367 (♂). Type-loc: Bettutan, nr Sandakan, N Borneo.
- nigroaenescens* Rondani. — Borneo, Java, Malaya, Sumatra.
1875, Annali Mus. civ. Stor. nat. Giacomo Doria 7: 422 (♂). Type-loc: Sarawak, Borneo.
- strigata* de Meijere, 1914, Tijdschr. Ent. 57: 142 (♀). Type-loc: Nongkodjadjar, Gunung Gede, Java.
- striga*, error.
- petulans* Curran, 1928, J. fed. Malay St. Mus. 14: 282 (♂ ♀). Type-locs: Ginting Sempak, Selangor-Pahang border; Jor Camp (1800 ft), Batang Padang, Perak; No. 4 camp (4800 ft), Cameron Highlands and Sungai Ringlet (3500 ft), Pahang, Malaya.
- nursci* Brunetti. — India [Himachal Pradesh].
1923, Fauna Br. India, Dipt. 3: 240 (♂). Type-loc: Simla, India.

ornata (Brunetti). — India [Uttar Pradesh].

1915, Rec. Indian Mus. 11: 232 (♂; *Merodon*). Type-loc: Bhowali, Ku-maon District, India.

pendleburyi Curran. — Borneo, Malaya.

1928, J. fed. Malay St. Mus. 14: 279 (♂). Type-loc: Kuala Tahan (300 ft), Pahang, Malaya.

penicillata Brunetti. — India [Assam].

1923, Fauna Br. India, Dipt. 3: 242 (♂). Type-loc: Lower Ranges, Khasi Hills, Assam, India.

spinipes Curran. — Malaya, Thailand.

1928, J. fed. Malay St. Mus. 14: 283 (♂). Type-loc: Khao Luang (3000 ft), Nakhon Si Thammarat, Thailand.

steyskali Thompson (for *spinipes* Shiraki). — Formosa. N. name.

spinipes Shiraki, 1930, Mem. Fac. Agric. Taihoku imp. Univ. 1: 72 (♂ ♀; *Zelima*; preocc. Curran 1928). Type-locs: Shinchiku, Karenko, Koshun, and Taiheizan, Formosa.

stylata Hull. — Java.

1944, Proc. ent. Soc. Wash. 46: 46 (♂). Type-loc: Soekaboemi, Java.

Subfamily MICRODONTINAE

Genus INDASCIA Keiser

Indascia Keiser, 1958, Revue suisse Zool. 65: 221. Type-species, *Ascia brachystoma* Wiedemann, by original designation.

brachystoma (Wiedemann). — ?India.

1824, Analecta Ent.: 33 (♀; *Ascia*). Type-loc: "India orientalis."

gracilis Keiser. — Ceylon.

1958, Revue suisse Zool. 65: 223 (♂). Type-loc: Bot. Gardens, Peradeniya, Ceylon.

Genus MICRODON Meigen

Microdon Meigen, 1803, Magazin InsektKde 2: 275. Type-species, *Musca mutabilis*

Linnaeus (as *Mulio mutabilis* Fabricius), by monotypy.

References: de Meijere 1908; Curran 1942, 1947 (keys).

Subgenus MICRODON Meigen

aenoviridis Curran. — Borneo, Malaya.

1931, J. fed. Malay St. Mus. 16: 302 (♂). Type-loc: Kedah Peak (1000-2000 ft), Malaya.

aenus Keiser. — Sumbawa.

1952, Verh. naturf. Ges. Basel 63: 172 (♂). Type-loc: Pogobina, W Sumbawa.

albofascia Hull. — Burma.

1944, Ann. Mag. nat. Hist. (11) 11: 253 (♀). Type-loc: Adung Valley, Burma.

Syrphidae

alboscutatus Curran. — Borneo, Malaya, Thailand.

1931, J. fed. Malay St. Mus. 16: 303 (♂ ♀). Type-loc: Langkawi Island, W Coast, Malaya.

annandalei Brunetti. — India [Assam, Himachal Pradesh], E & W Pakistan, Nepal.

1907, Rec. Indian Mus. 1: pl. xiii, fig. 10; 1908, Rec. Indian Mus. 2: 91 (♂). Type-loc: Soondrial, Nepal.

apidiformis Brunetti. — India [Assam].

1925, Rec. Indian Mus. 27: 78 (n. name for *apiformis* Brunetti).

apiformis Brunetti, 1923, Fauna Br. India, Dipt. 3: 314 (♂; preocc. De Geer 1776). Type-loc: Tura, Garo Hills, Assam, India.

auricinctus Brunetti. — Ceylon, Formosa, India [W Bengal], Philippines [Luzon].

1908, Rec. Indian Mus. 2: 93 (♀). Type-loc: Kandy, Ceylon.

aurilinea Hull. — Java.

1944, Ann. Mag. nat. Hist. (11) 11: 258 (♂ ♀). Type-loc: Soekaboemi, Java.

aurivesta Hull. — Java.

1950, Ann. Mag. nat. Hist. (12) 3: 611 (♂). Type-loc: Soekaboemi, Java.

auroscutatus Curran. — Malaya, Thailand.

1928, J. fed. Malay St. Mus. 14: 152 (♂ ♀). Type-locs: Kuala Tahan (300 ft) and Kuala Tekv (500 ft), Pahang, Malaya; and Khao Ram (300-750 ft), Nakhon Si Thammarat, Thailand.

var. *variventris* Curran. — Malaya.

1928, J. fed. Malay St. Mus. 14: 154 (♀). Type-loc: Kuala Tahan (300 ft), Pahang, Malaya.

baramus Curran. — Borneo.

1942, Am. Mus. Novit. 1216: 3 (♀). Type-loc: Baram, Sarawak, Borneo.

beatus Curran. — Borneo.

1942, Am. Mus. Novit. 1216: 4 (♂). Type-loc: Matang Road, Sarawak, Borneo.

bellus Brunetti. — India [Uttar Pradesh], Nepal.

1923, Fauna Br. India, Dipt. 3: 315 (♀). Type-loc: Mussoorie, India.

bicolor Sack. — Formosa.

1922, Arch. Naturgesch. (A) 87(11): 272 (♀). Type-loc: Anping, Formosa.

bicoloratus Hull. — Sumbawa.

1944, Ann. Mag. nat. Hist. (11) 11: 254 (♂). Type-loc: Sumbawa, Dutch East Indies (Indonesia).

caeruleus Brunetti. — Formosa, India [Assam].

1908, Rec. Indian Mus. 2: 92 (♀). Type-loc: Margherita, Assam, India.

var. *simplex* Shiraki. — Formosa; Japan.

1930, Mem. Fac. Agric. Taihoku imp. Univ. 1: 15 (♂ ♀). Type-locs: Japan-Yamaguchi; and Formosa (Asahi V).

carbonarius Brunetti. — India [Assam].

1923, Fauna Br. India, Dipt. 3: 314 (♂ ♀). Type-loc: Tura, Garo Hills, Assam, India.

chapini Hull. — Thailand.

1941, J. Wash. Acad. Sci. 31: 438 (♂). Type-loc: Patmeung Mts, Siam (Thailand).

- clavicornis* Sack. — Philippines [Luzon, Mindanao].
1926, Philipp. J. Sci. 29: 592 (♀). Type-loc: Paete, Laguna, Luzon, Philippines.
- contractus* Brunetti. — India [Maharashtra].
1923, Fauna Br. India, Dipt. 3: 310 (♀). Type-loc: Deesa, Bombay Presidency, India.
- conveniens* Brunetti. — India [Assam].
1923, Fauna Br. India, Dipt. 3: 311 (♀). Type-loc: Assam, India.
- dimidiatus* Curran. — Borneo.
1942, Am. Mus. Novit. 1216: 3 (♂). Type-loc: Lundu, Sarawak, Borneo.
- elisabeth* Keiser. — Ceylon.
1958, Revue suisse Zool. 65 (1): 211 (♂). Type-loc: Kandy, Lady Horton's, Ceylon.
- flavipes* Brunetti. — Burma.
1908, Rec. Indian Mus. 2: 92 (♂ ♀). Type-loc: Mergui, Lower Burma.
- formosanus* Shiraki. — Formosa.
1930, Mem. Fac. Agric. Taihoku imp. Univ. 1: 22 (♂ ♀). Type-locs: Horisha, and Masha, Formosa.
- fulvipes* de Meijere. — Sumatra, Malaya.
1908, Tijdschr. Ent. 51: 203 (♀). Type-loc: Tandjong, Sumatra.
- fulvopubescent* Brunetti. — Ceylon.
1923, Fauna Br. India, Dipt. 3: 313 (♀). Type-loc: Ceylon.
- fumipennis* Hull. — Java.
1944, Ann. Mag. nat. Hist. (11) 11: 259 (♂). Type-loc: Soekaboemi, Java.
- fuscicornis* Sasakawa. — Thailand.
1960, Nature Life S.E. Asia 1: 451 (♂). Type-loc: Doi Inthanon, Thailand.
- fuscus* de Meijere. — Sumatra.
1908, Tijdschr. Ent. 51: 204 (♀). Type-loc: Median, Sumatra.
- gloriosus* Hull. — Thailand.
1941, J. Wash. Acad. Sci. 31: 439 (♂). Type-loc: Patmeung Mts, Siam (Thailand).
- grandis* Curran. — Malaya.
1928, J. fed. Malay St. Mus. 14: 159 (♂). Type-loc: Jor Camp, Perak, F.M.S.
- indicus* (Dobeschall). — Bali: Moluccas (Amboina).
1857, Natuurk. Tijdschr. Ned.-Indië 14: 404 (sex?; *Ceratophya*). Type-loc: Amboina, Moluccas. Reference: de Meijere 1908: 211 (redescr.).
- investigator* Hull. — Philippines [Mindanao].
1937, Psyche, Camb. 44: 20 (♂). Type-loc: Galog River, Mt Apo, Davao, Mindanao, Philippines.
- klossi* Curran. — Borneo.
1931, J. fed. Malay St. Mus. 16: 343 (♂). Type-loc: Bettotan, nr Sandakan, N Borneo.
- lanka* Keiser. — Ceylon.
1958, Revue suisse Zool. 65: 213 (♀). Type-loc: Carney, Ratnapura, Ceylon.
- latiscutellaris* Curran. — Borneo.
1931, J. fed. Malay St. Mus. 16: 341 (♀). Type-loc: Bettotan, nr Sandakan, N Borneo.

- lativentris* de Meijere. — Sumatra.
1921, Tijdschr. Ent. 64: lii; also 1924, Tijdschr. Ent. 67 (Suppl.): 19 (♂). Type-loc: Aur Kumandis, Sumatra.
- laxiceps* Curran. — Borneo.
1942, Am. Mus. Novit. 1216: 2 (♀). Type-loc: Trusan, Sarawak, Borneo.
- lundura* Curran. — Borneo.
1942, Am. Mus. Novit. 1216: 3 (♂). Type-loc: Lundu, Sarawak, Borneo.
- luxor* Curran. — Malaya.
1931, J. fed. Malay St. Mus. 16: 306 (♂). Type-loc: Bukit Kutu (3500 ft), Selangor, Malaya.
- metallicus* de Meijere. — India [W Bengal].
1904, Bijdr. Dierk. 18: 98 (♂). Type-loc: Darjeeling, India.
- minuticornis* Curran. — Borneo.
1931, J. fed. Malay St. Mus. 16: 342 (♂ ♀). Type-loc: Bettotan, nr Sandakan, N Borneo.
- montis* Keiser. — Ceylon.
1958, Revue suisse Zool. 65: 214 (♂). Type-loc: Pidratalagala, Ceylon.
- pagdeni* Curran. — Malaya.
1942, Am. Mus. Novit. 1216: 6 (♀). Type-loc: Kedah Peak, Kedah (3284 ft), Malaya.
- pendelburyi* Curran. — Malaya.
1931, J. fed. Malay St. Mus. 16: 305 (♀). Type-loc: Kedah Peak (3000–3500 ft), Malaya.
- pretiosus* Curran. — Malaya.
1931, J. fed. Malay St. Mus. 16: 304 (♂). Type-loc: Langkawi Island, W coast Malay Peninsula, Malaya.
- robinsoni* Curran. — Malaya, Thailand.
1928, J. fed. Malay St. Mus. 14: 154 (♀). Type-loc: Nakhon Si Thammarat, Khao Luang (2000 ft), Thailand.
- ruficaudus* Brunetti. — Formosa, India [W Bengal].
1907, Rec. Indian Mus. 1: pl. xiii. fig. 11; 1908, Rec. Indian Mus. 2: 93 (♀). Type-loc: Calcutta, India.
- rutiliventris* Vockeroth (for *rufiventris* Curran). — Borneo. N. name.
rufiventris Curran, 1942, Am. Mus. Novit. 1216: 4 (♀; preocc. Rondani 1848). Type-loc: Kuching, Sarawak, Borneo.
- rutilus* Keiser. — Sumbawa.
1952, Verh. naturf. Ges. Basel 63: 173 (♂). Type-loc: Pogobina, W Sumbawa.
- sacki* van der Goot. — Philippines [Mindanao].
1964, Beaufortia 10: 220 (n. name for *flavipennis* Sack).
- flavipennis* Sack, 1926, Philipp. J. Sci. 29: 593 (♂; preocc. Curran 1925). Type-loc: Surigao, Mindanao, Philippines.
- var. *nigrita* Hull. — Philippines [Mindanao].
1944, Ann. Mag. nat. Hist. (11) 11: 257 (♂; *flavipennis* var.). Type-loc: Mindanao, Philippines.
- simplicicornis* de Meijere. — Java.
1908, Tijdschr. Ent. 51: 205 (♂). Type-loc: Buitenzorg (Bogor), Java.
- squamipennis* Brunetti. — India [Assam].
1923, Fauna Br. India, Dipt. 3: 316 (♀). Type-loc: Tura, Garo Hills, Assam, India.

- stenogaster* Curran. — Borneo.
1931, J. fed. Malay St. Mus. 16: 344 (♂ ♀). Type-loc: Bettutan, nr Sandakan, N Borneo.
- stilboides* Walker. — Forniosa, India [W Bengal], Java, Philippines [Mindanao].
1849, List Dipt. Colln Br. Mus. 3: 538 (♀). Type-loc: East Indies.
- sulcatus* Hull. — Java.
1944, Ann. Mag. nat. Hist. (11) 11: 256 (♀). Type-loc: Soekaboemi, Java.
- sumatranaus* van der Wulp. — Sumatra.
1881, Midden-Sumatra Exped., Dipt. 4: 29 (♀). Type-loc: "Loeboek gedang," Sumatra.
- sumbanus* Keiser. — Sumbawa.
1952, Verh. naturf. Ges. Basel 63: 174 (♀). Type-loc: Pogobina, West Sumbawa.
- taprobanicus* Keiser. — Ceylon.
1958, Revue suisse Zool. 65: 212 (♂). Type-loc: Kandy, Hantana, Ceylon.
- tricinctus* de Meijere. — Java, Philippines [Mindanao].
1908, Tijdschr. Ent. 51: 208 (♂ ♀). Type-loc: Batavia (Djakarta), Java.
- trigonospilus* Bezzii. — China [Kwangtung].
1927, Bull. Lab. Zool. Portici. 20: 4 (♀). Type-loc: La Kow, Honan (Canton), China.
- trimacula* Curran. — Malaya.
1928, J. fed. Malay St. 14: 156 (♂). Type-locs: Perak, Batang, Padang, Tapah-Pahang Rd (12th mi), and Jor Camp, Malaya.
- unicolor* Brunetti. — India [Orissa].
1915, Rec. Indian Mus. 11: 255 (♂). Type-loc: Puri, Orissa, India.
- varicornis* Sack. — Philippines [Mindanao].
1926, Philipp. J. Sci. 29: 594 (♀). Type-loc: Butuan, Agusan, Mindanao, Philippines.
- vespiformis* de Meijere. — Java, Malaya.
1908, Tijdschr. Ent. 51: 210 (sex?). Type-loc: Java.
- wulpiae* Mik. — Sumatra.
1899, Wien. ent. Ztg 18: 143 (n. name for *apicalis* van der Wulp).
apicalis van der Wulp, 1881, Midden-Sumatra Exped., Dipt. 4: 29 (♀; preocc. Walker 1859). Type-loc: Koetoer, Sumatra.

Subgenus ARCHIMICRODON Hull

- Microdon*, subg. *Archimicrodon* Hull, 1945, Proc. New Engl. zool. Club 23: 75.
Type-species, *digitator* Hull, by original designation.
- digitator* Hull. — Java.
1937, Psyche, Camb. 44: 19 (♂). Type-loc: Tjibodas, Mt Gede, Java.

Subgenus KRYPTOPYGA Hull

- Kryptopyga* Hull, 1944, J. Wash. Acad. Sci. 34: 129. Type-species, *pendulosa* Hull, by original designation.

pendulosus Hull. — Java.

1944, J. Wash. Acad. Sci. 34: 130 (♂). Type-loc: Soekaboemi, Java.

Genus PARAMICRODON de Meijere

Paramicrodon de Meijere, 1913, Nova Guinea 9: 360. Type-species, *lorenzii* de Meijere, by monotypy.

Syrphinella Hervé-Bazin, 1926, Encycl. ent. (B) II, Dipt. 3: 73. Type-species, *miranda* Hervé-Bazin, by monotypy.

Myxogasteroides Shiraki, 1930, Mem. Fac. Agric. Taihoku imp. Univ. 1: 9. Type-species, *Myxogaster nigripennis* Sack, by original designation. N. synonym.

cinetellus (Sack). — Philippines [Mindanao].

1926, Philipp. J. Sci. 29: 590 (♀; *Myxogaster*). Type-loc: Agusan, Butuan, Mindanao, Philippines.

dicipliens de Meijere. — Java.

1917, Tijdschr. Ent. 60: 242 (♀). Type-loc: Salatiga, Java.

mirandus (Hervé-Bazin). — Indochina.

1926, Encycl. ent. (B) II, Dipt. 3: 74 (♀; *Syrphinella*). Type-loc: Chuachan, Indochina.

nigripennis (Sack). — Formosa.

1922, Arch. Naturgesch. (A) 87(11): 275 (♂; *Myxogaster*). Type-loc: Toa Tsai Kutsu, Formosa. N. comb.

novus Hull. — Philippines [Mindanao].

1937, Psyche, Camb. 44: 22 (♀). Type-loc: Galog River (6500 ft), Mt Apo, Davao, Mindanao, Philippines.

Microdon (Genus PAROCYPTAMUS) Shiraki

Parocryptamus Shiraki, 1930, Mem. Fac. Agric. Taihoku imp. Univ. 1: 11. Type-species, *sonamii* Shiraki, by original designation.

Stenomicodon Hull, 1937, Psyche, Camb. 44: 26. Type-species, *purpureus* Hull, by original designation.

purpureus (Hull). — Formosa.

1937, Psyche, Camb. 44: 26 (♀; *Stenomicodon*). Type-loc: Tainan, Formosa.

sonamii Shiraki. — Formosa.

1930, Mem. Fac. Agric. Taihoku imp. Univ. 1: 12 (♂). Type-locs: Shinchiku, Sokutsu, and Chipon, Formosa.

Genus PARAMIXOGASTER Brunetti

Paramixogaster Brunetti, 1923, Fauna Br. India, Dipt. 3: 320. Type-species, *Mixogaster vespiformis* Brunetti, by original designation.

icariiformis Pendlebury. — Malaya.

1927, Proc. ent. Soc. Lond. 1: 38 (♀). Type-loc: Bukit Kuta (3500 ft), Selangor, Malaya.

vespiformis (Brunetti). — India [Assam].

1913, Rec. Indian Mus. 8: 169 (♂; *Mixogaster*). Type-loc: Dibrugarh, Assam, India.

wegneri Keiser. — Moluccas (Amboina).

1964, Mitt. schweiz. ent. Ges. 37: 84 (♂ ♀). Type-loc: Ambo Island, Indonesia.

Genus PARAMIXOGASTEROIDES Shiraki

Paramixogasteroides Shiraki, 1930, Mem. Fac. Agric. Taihoku imp. Univ. 1: 8.

Type-species, *Myxogaster variegata* Sack, by monotypy.

variegata (Sack). — Formosa.

1922, Arch. Naturgesch. (A) 87(11): 274 (♀; *Myxogaster*). Type-loc: Toa Teai Kutau, Formosa.

Unplaced genus and species of Syrphidae

Genus PRIOMERUS Macquart *Palumbis* (Korinko)

Priomerus Macquart, 1834, Hist. nat. Ins. Dipt. 1: 511. Type-species, *fasciatus* Macquart, by monotypy.

fasciatus Macquart. — India.

1834, Hist. nat. Ins. Dipt. 1: 511 (♂). Type-loc: East India.

Family CONOPIDAE

By Kenneth G. V. Smith

Department of Entomology, British Museum (Natural History),
London, England.

The Conopidae are a relatively small family of parasitic flies well represented in the major zoogeographical regions. Most are parasitic on aculeate Hymenoptera and are usually found among flowers in dry areas where their hosts abound. The Stylogasterinae are parasitic on cockroaches and calyptrate Diptera. Many species appear to mimic wasps among which they occur, although some are parasitic on bees of quite different appearance.

General keys to the Conopidae of the Oriental region are given by Brunetti (1923), Chen (1939) and Kröber (1939–40), and other useful keys are included in Camras (1957a, *metaxanthus* group of *Conops*), Chvála (1953, *Sicus*), and Smith (1960, world *Pleurocerinella*; 1966a, Indo-Australian *Stylogaster*). The biology of Conopidae is reviewed by Smith (1966b, 1967).

Subfamily CONOPINAE

Genus ABRACHYGLOSSUM Kröber

Abrachyglossum Kröber, 1917, Arch. Naturgesch. (A) 83(7): 142. Type-species, *Conops capitatum* Loew, by original designation.

wui Ōuchi. — China [Chekiang].

1939, J. Shanghai Sci. Inst. (3) 4: 195 (♂). Type-loc: Tienmushan, Chekiang Province, China.

Genus ARCHICONOPS Kröber

Archiconops Kröber, 1939, Ann. Mag. nat. Hist. (11) 4: 381. Type-species, *Conops insularis* Kröber, by present designation.

erythrocephala (Fabricius). — Burma, Celebes, Ceylon, China [Kiangsu], India [Madras, Maharashtra, Sikkim, W Bengal], Indochina; Africa.

1794, Ent. Syst. 4: 392 (sex?; *Conops*). Type-locs: India orientali; Africa. morosa (de Meijere). — Java.

1924, Tijdschr. Ent. 76: 213 (♂ ♀; *Conops*). Type-loc: Kloet, ca 300 m, Java.

Genus CONOPS Linnaeus

Conops Linnaeus, 1758, Syst. Nat. Ed. 10: 640. Type-species, *flavipes* Linnaeus, by designation of Curtis (1831; pl. 377).

Subgenus ASICONOPS Chen

Conops, subg. *Asiconops* Chen, 1939, Notes Ent. chin. 6(10): 171. Type-species, *aureomaculatus* Kröber, by original designation.